

---

# Wet and dry weather toxicity in the San Gabriel River

---

Kenneth Schiff, Beth Bax<sup>1</sup>, Phil Markle<sup>1</sup>, Terry Fleming<sup>2</sup> and Jennifer Newman<sup>3</sup>

## ABSTRACT

The lower San Gabriel River is an urban watershed located on the border of Los Angeles and Orange Counties. It has a diversity of potential pollutant sources including five water reclamation plants (WRPs) that discharge treated wastewaters and more than 100 storm drains that discharge largely untreated urban runoff to the river. The goal of this study was to assess the magnitude of toxicity to *Ceriodaphnia dubia* throughout the lower San Gabriel River watershed during wet and dry weather, identify the responsible toxicants, and compare the magnitude of toxicity over time to evaluate the effectiveness of previous watershed management actions. Wet weather runoff was sampled from sites located at the end of the four main reaches of the lower San Gabriel River; Walnut Creek, San Jose Creek, Coyote Creek, and San Gabriel River mainstem. None of the samples collected over two wet seasons exhibited acute or chronic toxicity. Dry weather samples were tested from 16 locations distributed throughout the lower watershed for up to 18 months. None of the dry weather samples from Walnut Creek, San Jose Creek, or the San Gabriel River mainstem exhibited acute or chronic toxicity. Acute and chronic toxicity was intermittently measured in the Coyote Creek tributary. Toxicity identification evaluations suggested nonpolar organic constituents, likely diazinon and perhaps surfactants, as possible toxicants. Toxicity observed in this study was significantly reduced compared to a similar study of the watershed 12 years previously, especially in the San Gabriel River mainstem. Much of the reduction in toxicity was associated with upgrades in WRP treatment. Little to no change in toxicity was observed in Coyote Creek upstream of the WRP discharge where little to no control of dry weather urban runoff had occurred.

## INTRODUCTION

Urban watersheds receive a multitude of potential pollutants that can affect aquatic life (Bay *et al.* 1996, Ackerman *et al.* 2005, Tiefenthaler and Stein 2005). The San Gabriel River, located on the border between Los Angeles and Orange Counties in southern California, is an ideal example of the ways in which aquatic life may be impacted by potential pollutants. Sources of potential pollutants include treated sanitary wastewaters from five Water Reclamation Plants (WRPs) and untreated urban runoff from approximately 350 km<sup>2</sup> of developed land discharged into the river via a municipal separate storm sewer system, as well as once-through cooling waters from two power generating stations that is mixed with low volume industrial and sanitary wastes then discharged into the watershed's estuary.

To complicate the fate and transport of anthropogenic pollutants and their resultant effects on aquatic life, the hydrology of many urban watersheds is often highly modified. For example, three major dams were constructed in the upper undeveloped reaches of the San Gabriel River watershed in order to capture, retain, and utilize wet season runoff for potable water use during the dry season. While this provides much needed water for the citizens of Los Angeles, the upper watershed is now hydrologically disconnected from the urbanized lower watershed. The result is that natural waters are unavailable for mixing and dispersion of anthropogenic discharges downstream. Even greater hydromodification exists in the urbanized lower San Gabriel River watershed. Many miles of the river in this portion of the watershed are lined with concrete in an effort to reduce flooding and property damage, but this modification also results in the maximum exposure of pollutants to aquatic life through the loss of natural stream and treatment processes. Where unlined channels exist in the lower water-

---

<sup>1</sup> Los Angeles County Sanitation Districts, Whittier, CA

<sup>2</sup> US Environmental Protection Agency, San Francisco, CA

<sup>3</sup> Los Angeles Regional Water Control Board, Los Angeles, CA

shed, temporary dams are inflated to enhance groundwater recharge.


In response to pollutant inputs and hydrologic modification, many urban watersheds have been the focus of water quality regulatory efforts. Urban Los Angeles once again provides a good example. More than 180 waterbodies in the Los Angeles region have been placed on the United States Environmental Protection Agency's (EPA's) list of impaired waters. This list, also referred to as the 303(d) list (referring to section 303d of the Clean Water Act), identifies locations impacted by specific pollutants that can result in toxicity to aquatic life and other impacts. The effect of the 303(d) list is the mandate for future regulation (termed a total maximum daily load or TMDL), which will require the mitigation of these pollutant inputs.

In the San Gabriel River watershed, managers have been implementing mitigation to negate the effects of these pollutant inputs. Over the past 10 years, WRPs in the San Gabriel River watershed have installed additional treatment processes, costing over \$40 million, which have dramatically improved the water quality of their discharges. Controlling pollutant impacts due to urban runoff has been more difficult. Up to \$10 million has been spent on structural best management practices (BMPs) in the San Gabriel River, yet few (if any) trends in concentrations of toxic constituents monitored have been observed (LACDPW 2005). Unlike WRPs, urban runoff discharges are diffuse and, as a result, perhaps more difficult to treat and/or control.

The objective of this study was to evaluate the impact of pollutants on aquatic life in the highly urbanized lower watershed of the San Gabriel River. Impact to aquatic life was assessed through the use of toxicity testing. Four specific goals were identified: 1) assess the magnitude of toxicity at selected locations throughout the San Gabriel River watershed; 2) determine whether or not this magnitude changes seasonally; 3) if toxicity exists, identify the responsible toxicants; and 4) compare the magnitude of toxicity in this study to studies conducted historically in the San Gabriel River watershed to evaluate the effectiveness of watershed management actions.

## METHODS

Toxicity in the San Gabriel River watershed was evaluated by separating the study into wet weather and dry weather components (Figure 1; Table 1).



**Figure 1. Map of the lower San Gabriel River watershed including dry and wet weather sampling locations.**

The wet weather component consisted of four sampling sites located at the downstream end of major reaches that receive urban runoff. Twenty-liter flow weighted composites were sampled during three storm events on December 29, 2004 (5.3 cm precipitation), April 22, 2005 (2.2 cm precipitation), and January 1, 2006 (3.7 cm precipitation). The dry weather component consisted of sampling a total of 10 sites that included the same four sites sampled during wet weather, plus an additional six sites strategically located in the immediate vicinity of WRP discharges or urban runoff inputs. Dry weather samples were collected at least three days after rain events. Twenty-liter samples were collected from each site during dry weather on a monthly basis from March 2005 to February 2006. Within seven months of this study's initiation, an additional six sites were added for dry weather sampling, all in a single tributary (North Coyote Creek), as a result of observed toxicity. All sites from the Coyote Creek subwatershed, including the additional sites in North Coyote Creek, were sampled until August 2006.

**Table 1. Station Location Information (NAD83 datum).**

Site	Water Body	Location	Latitude/ Longitude
1	Walnut Creek	Walnut Creek At Merced Ave	N34° 03'53.1" W117° 57'09.6"
2	Walnut Creek	At Baldwin Park Blvd	N34° 03'47.7" W117° 58'54.5"
3	San Jose Creek Reach 1	San Jose Creek at access gate from SJCWRP/JAO – Upstream of SJCWRP, approximately 100 yards downstream of Workman Mill Rd	N34° 02'06.7" W118° 01'14.9"
4	San Gabriel River Reach 3	San Gabriel River at Peck Rd - Downstream of confluence of SGR with SJC	N34° 02'02.9" W118° 02'20.2"
5	San Gabriel River Reach 1	San Gabriel River at Spring St - Downstream of LCWRP outfall 001	N33° 48'38.9" W118° 05'26.8"
6	Coyote Creek	Coyote Creek at pedestrian foot bridge south of LBWRP - Downstream of LBWRP outfall 001 and upstream of estuary	N33° 47'41.9" W118° 05'22.0"
7	Coyote Creek	Coyote Creek at Cerritos Ave - Upstream of LBWRP outfall; downstream of entrance of Carbon Creek into Coyote Creek	N33° 48'36.9" W118° 04'33.3"
8	Coyote Creek	Coyote Creek at Centralia Ave - Downstream of confluence with Fullerton Creek and an industrial drain	N33° 50'19.3" W118° 03'37.9"
9	Coyote Creek	Coyote Creek at Artesia Blvd- Downstream of Brea Creek/Coyote Creek confluence	N33° 52'23.7" W118° 01'08.0"
10	Coyote Creek (North Fork)	North fork of Coyote Creek at Alondra Blvd- Downstream of La Mirada Creek	N33° 53'15.4" W118° 01'58.9"
11	Coyote Creek (North Fork)	Coyote Creek North Fork at La Mirada Creek/Coyote Creek confluence	N 33° 53.731' W 118° 02.155'
12	Coyote Creek (North Fork)	Coyote Creek North Fork- 1.0 mile upstream of Alondra	N 33° 54.133' W 118° 02.488'
13	Coyote Creek (North Fork)	Coyote Creek North Fork - 2.0 miles upstream of Alondra	N 33° 54.862' W 118_02.346'
14	Coyote Creek (North Fork)	Coyote Creek North Fork - 2.5 miles upstream of Alondra	N 33° 55.411' W 118° 02.138'
15	La Mirada Creek	La Mirada Creek before entering Coyote Creek North Fork	N 33° 53.503' W 118° 01.846'
16	Milan Creek	Milan Creek before entering Coyote Creek North Fork	N 33° 54.369' W 118° 02.422'

All samples were tested for toxicity using *Ceriodaphnia dubia* examining both acute (lethality) and chronic (reproductive success) endpoints. Testing was initiated within 36 hours of sample collection using undiluted sample and a negative control following standard EPA protocols (EPA 2003a; Table 2). Test organisms were obtained from in-house brood cultures and test duration/exposure lasted until 60% of the surviving females in the control had released three broods (typically between six and seven days). Test solutions were renewed daily.

Toxicity was defined as a 25%, or greater, organism response in the sample exposure relative to control organism response (i.e., <75% survival or reproduction in the 100% sample exposure). In addition, hypothesis testing was conducted following EPA guidelines (EPA 2003a). Hypothesis testing consisted of the nonparametric Fisher's Exact Test for the

survival endpoint and an analysis of variance (ANOVA) followed by a multiple comparison procedure for the reproduction endpoint. The parametric Dunnett's Test was used to identify statistically significant differences from the control for reproduction data that were normally distributed with homogeneous variances. The nonparametric Steel's Many-One Test was employed when the data failed normal distribution or equality of variance assumptions.

If a sample was toxic, a toxicity identification evaluation (TIE) was initiated (USEPA 1991, 1993b). TIE testing used the remaining sample, stored at 4°C, within seven days of baseline test conclusion. For those samples in which only the reproductive endpoint elicited a toxic response, only 100% and control concentrations were evaluated in the TIE. In these cases, the TIE consisted of a full seven-day chronic test with each sample manipula-

**Table 2. Test conditions and requirements.**

Test Organism	<i>Ceriodaphnia dubia</i>
Organism Source:	In-house Cultures
Organism Age at Initiation:	<24 hours old and released within an eight hour period
Test Duration:	Until 60% or ore of the surviving females have three broods
Concentrations Tested:	0% and 100%
Solution Renewal:	Daily
Feeding:	0.1 ml YCT and 0.1 <i>Selenastrum</i> algal suspension daily
Test Chamber:	50 ml Disposable
Solution Volume:	15 ml
Control Water:	Either diluted mineral water (8 parts deionized water: 2 parts Perrier® water) or Reconstituted deionized water (hard)
Number of Replicates:	10
Organisms per Replicate:	1 assigned by blocking by known parentage
Photoperiod:	16 hours light (50 - 100 ft-c), 8 hours dark
Test Temperature:	25 + 1°C.
Endpoints Measured:	Survival and Reproduction
Test Acceptability Criteria:	80% or greater survival with an average of 15 or more young per surviving female in the control organisms. 60% of surviving females in the controls must produce three broods within 8 days.

tion consisting of 10 replicates, with daily renewals. For those samples where the survival endpoint elicited a toxic response, three dilutions (25, 50, 100%) and a control were evaluated using four replicates containing five test organisms each. In the case of a TIE in response to survival, the exposure duration was 96 hours, with renewal after 48 hours.

The TIE manipulations focused on both characterization and identification phases (EPA 1991, 1993b). These manipulations included: 1) pH adjustment; 2) aeration; 3) Ethylenedinitrilo-Tetraacetic Acid (EDTA); 4) Sodium thiosulfate (STS); 5) filtration; 6) piperonyl butoxide (PBO); 7) anion exchange column; 8) solid phase extraction (SPE); 9) SPE elution; and 10) no manipulation. By conducting each of these manipulations, the results, alone or in combination, can help to identify the responsible toxicant(s) (Table 3).

All quality assurance/quality control criteria were met for this study. These criteria included all of the test acceptability criteria (Table 2). In addition, positive control samples using reference toxicants (copper chloride) confirmed the relative sensitivity and stability of test organisms during the course of the study.

## RESULTS

None of the storms sampled during this study were acutely or chronically toxic to *C. dubia*. At all four sites, during all three storms, survival and reproduction were greater than 75% relative to controls.

Eighteen of 196 (9%) total dry weather samples exhibited chronic toxicity during this study (Table 4). Twelve of 196 (6%) total dry weather samples exhibited acute toxicity during this study. All of the dry weather samples that exhibited acute toxicity also exhibited chronic toxicity. In only one case was statistically significant toxicity observed when the response was less than 25% relative to controls (Station 15, Jan 2006). Only once was toxicity greater than 25% relative to controls and not statistically significant (Station 15, Mar 2006).

All observed toxicity during this study was from Coyote Creek (Table 4). No toxicity was observed in Walnut Creek, San Jose Creek, or San Gabriel River Reaches 1 or 3. Widespread toxicity in

**Table 3. Toxicity Identification Evaluation (TIE) sample manipulations and their respective interpretations.**

TIE Sample Manipulation	Expected response
pH Adjustment (pH 7 and 8.5)	Alters toxicity in pH sensitive compounds (i.e., ammonia and some trace metals)
Aeration	Reduces toxicity attributable to volatile, sublutable, and/or easily oxidizable compounds
Ethylenedinitrilo-Tetraacetic Acid (EDTA) Addition	Chelates trace metals, particularly divalent, cationic metals
Sodium thiosulfate (STS) Addition	Reduces toxicants attributable to oxidants (i.e., chlorine) and some trace metals
Filtration	Removes toxicity related to and/or associated with particulates
Solid Phase Extraction (SPE) with C <sub>18</sub>	Removes toxicity associated with non-polar organics (i.e., pesticides, surfactants)
Sequential Solvent Extraction of with C <sub>18</sub> Column	SPE extraction can be used to confirm toxicity due to nonpolar organic compounds. Sequential extraction using solvents of gradually decreasing polarity can separate these compounds into fractions providing further toxicant resolution and isolation for chemical analysis
Piperonyl Butoxide (PBO)	Removes toxicity caused by metabolically activated pesticides (i.e., organophosphorous pesticides). Increases toxicity attributable to pyrethroid pesticides
Anion Exchange	Removes toxicity associated with anionic compounds, including some trace metals and surfactants
No Manipulation	For comparing the relative effectiveness of other manipulations and quantifies the persistence of toxicity in the stored sample



**Table 4. Summary of dry weather *Ceriodaphnia dubia* toxicity from San Gabriel River from March 2005 through August 2006.**

Location	Month of Sample Collection																		
	Mar. 2005	Apr. 2005	May 2005	Jun. 2005	Jul. 2005	Aug. 2005	Sept. 2005	Oct. 2005	Nov. 2005	Dec. 2005	Jan. 2006	Feb. 2006	Mar. 2006	Apr. 2006	May 2006	Jun. 2006	Jul. 2006	Aug. 2006	
<b>Walnut Creek</b>																			
1	-	-	-	-	-	-	-	-	-	-	-	-	-	-					
2	-	-	-	-	-	-	-	-	-	-	-	-	-	-					
<b>San Jose Creek Reach 1</b>																			
3	-	-	-	-	-	-	-	-	-	-	-	-	-						
<b>San Gabriel River Reach 3</b>																			
4	-	-	-	-	-	-	-	-	-	-	-	-	-						
<b>San Gabriel River Reach 1</b>																			
5	-	-	-	-	-	-	-	-	-	-	-	-	-						
<b>Coyote Creek</b>																			
6	-	S <sup>1</sup>	-	-	-	L <sup>1</sup> S <sup>1</sup>	-	-	-	-	-	-	-						
7	-	L <sup>1</sup> S <sup>1</sup>	-	-	-	L <sup>1</sup> S <sup>1</sup>	-	-	-	-	-	-	-						
8	-	L <sup>1</sup> S <sup>1</sup>	-	S <sup>1</sup>	-	L <sup>1</sup> S <sup>1</sup>	-	-	-	-	-	-	-						
9	S <sup>1</sup>	-	-	-	-	-	L <sup>1</sup> S <sup>1</sup>	-	-	-	-	-	-	-	-	-	-	-	-
10	S <sup>1</sup>	L <sup>1</sup> S <sup>1</sup>	-	L <sup>1</sup> S <sup>1</sup>	-	L <sup>1</sup> S <sup>1</sup>	-	-	-	-	-	-	-	-	-	-	-	-	-
11						-	S <sup>1</sup>	-	-	-	-	-	-	-	-	-	-	-	-
12						-	L <sup>1</sup> S <sup>1</sup>	-	-	-	-	-	-	-	-	-	-	-	-
13						-	L <sup>1</sup> S <sup>1</sup>	-	-	-	-	-	-	-	-	-	-	-	-
14						-	L <sup>1</sup> S <sup>1</sup>	-	-	-	-	-	-	-	-	-	-	-	-
15								-	-	-	-	-1	-	S	-	-	-	-	-
16								-	-	-	-	-	-	-	-	-	-	-	-


Shaded areas = samples not collected, "-" represents not toxic with effects less than 25% relative to control, L = Lethal, effect with toxicity less than 75% relative to control, S = Sub-lethal effect; reproduction less than 75% relative to control, "\*" = statistically significant from control

Coyote Creek was observed in April 2005. As a result, an additional six stations upstream were added between July and October 2005. Widespread toxicity was observed again in August 2005. Widespread toxicity was not observed again for the remaining 12 months (September 2005 to August 2006).

In the two events for which widespread toxicity was observed in Coyote Creek (April and August 2005), the toxicity appeared to originate in the upper portions of the tributary (Figure 2). In April 2005, 100% reproductive impairment was observed at the site sampled furthest upstream (Site 10) and reproductive success remained minimal moving downstream. *C. dubia* survival was also severely impacted at the furthest upstream station, then survival slowly increased downstream of the WRP discharge (Sites 7 and 6) indicating a potential dilution effect from the WRP effluent. The WRP in this reach was

discharging 13 mgd of effluent to Coyote Creek upstream of Site 6 during this sampling event. In August 2005, severe reproductive impairment was again observed at the site sampled furthest upstream (Site 14) and reproductive success remained minimal moving downstream. The WRP in this reach was not discharging effluent to Coyote Creek during this sampling event. *C. dubia* survival was more sporadic moving downstream during August 2005. Seventy eight percent survival was measured at Site 14 and decreased to 0% survival for downstream Sites 13 and 12. Survival increased to 100% at Site 11, but fell back to 0% survival for the remaining seven miles of Coyote Creek. The sudden increase in survival at Site 11 remains unexplained.

Seven TIEs were initiated during the study on dry weather samples exhibiting a 25% or greater effect (Table 5). Toxicity was no longer present for three of



**Figure 2. Survival in Coyote Creek April 2005 and August 2005.**

the samples (Sites 9 and 10 March 2005, Site 15 March 2006); consequently, no toxicant was identified.

Organophosphorus pesticides, most likely diazinon, were identified as the causative agent in one sample (Site 9 April 2005). This result was based on the exclusive removal of toxicity using SPE and the addition of PBO, which removes non-polar organic toxicants and inhibits toxicity due to diazinon, respectively (Figure 3). The SPE was sequentially eluted and these fractions were subsequently tested. Toxicity was recovered in the 80% methanol elution of the SPE column, a fraction associated with many organophosphorus pesticides including diazinon


(Figure 4). Finally, 1,700 µg/L diazinon was quantified in the sample using Enzyme-Linked Immunosorbent Assay (ELISA) techniques.

A non-polar organic toxicant(s), possibly a surfactant(s), was identified as the causative agent in the remaining three samples (Site 10 April, June, and August 2005). This result was based on the removal of toxicity using SPE. Toxicity was recovered in the 75% methanol elution, a fraction commonly associated with organophosphorus pesticides with surfactant toxicity recovery also documented (Norberg-King *et al.* 2005). An anion exchange column was used on two samples, with complete removal of toxicant

**Table 5. Summary of dry weather TIE results.**

Site	Sample Date	No Manipulation	TIE RESULTS (Survival in 100% Sample)										
			STS <sup>a</sup>	EDTA <sup>b</sup>	pH 7.0	pH 8.5	PBO <sup>c</sup>	Aeration	Filtration	Centrifuge	SPE	Anion	
9	Mar-05												
10	Mar-05												
10	Apr-05	0%	0%	0%	0%	0%	0% <sup>d</sup>	35%	0%	NT	87.5% <sup>e</sup>	NT	
10	Jun-05	0%	0%	0%	0%	0%	0%	10%	10%	30%	100% <sup>e</sup>	100%	
10	Aug-05	0%	0%	0%	0%	0%	0%	0%	0%	NT	100% <sup>e</sup>	0% <sup>f</sup>	
9	Sep-05	0%	0%	0%	0%	0%	100% <sup>g</sup>	0%	0%	NT	100% <sup>e</sup>	0%	
15	Mar-06												


NT = Not tested  
a – Sodium thiosulfate addition, two treatments of 10 and 25 ppm  
b – Ethylenedinitrio-tetraacetic acid addition, two treatments of 25 and 50 ppm  
c – Piperonyl butoxide addition, two treatments of 50 and 100 ppb  
d – 5% survival observed in the 50 ppb treatment with 0% survival in the 100 ppb treatment  
e – Toxicity recovered in only the 75% methanol elution  
f – Survival observed in lower concentrations of the sample indicating partial toxicity removal  
g – 80% survival observed in 50 ppb treatment and 100% survival in 100 ppb treatment



**Figure 3. Acute Phase I TIE - Site 9 sample collected on April 21, 2005.**

city observed in one sample (June 2005) and partial removal in the other (August 2005). This may be indicative of anionic surfactants, but might also suggest the presence of some trace metals. Elution of the anion column would help to confirm anionic surfactant toxicity, but attempts to recover toxicity from the anion column were not successful. However, other treatments to identify trace metals did not

reduce toxicity (i.e., EDTA), which helps to rule-out metals as a major source of toxicity. Aeration partially removed toxicity in the April 2005 sample. Some surfactants can be removed or partially removed through aeration. Finally, PBO did not reduce toxicity, and levels of diazinon in these three samples were low (<100 µg/L).



**Figure 4. Acute Phase I TIE Solid Phase Extraction Elution Testing - Site 9 sample collected on April 21, 2005.**

## DISCUSSION

Toxicity was not widespread in the San Gabriel River watershed over the 18 months examined during this study. No toxicity was observed at any site during any of the storm events sampled. Similarly, no toxicity was observed in four of the five major reaches in the lower watershed during dry weather. In Coyote Creek where toxicity was observed, the toxicity was intermittent and occurred only during 6 of the 18 sampling periods. This was despite an adaptive monitoring strategy, in which the number of sites sampled in Coyote Creek was doubled and the sampling period was extended by six months.


The lack of toxicity observed in this study was in direct contrast to historical studies in this watershed. While 9% of the samples were toxic in 2005/06, 55% of the samples collected for a similar study in 1992/93 were toxic (Bailey *et al.* 1995). Moreover, toxicity was observed in only a single reach (Coyote Creek) in 2005/06, while Bailey *et al.* (1995) identified toxicity in all five major reaches in the lower San Gabriel River watershed.

The difference in toxicity from tests conducted 14 years ago is likely due to changes in water quality. Bailey *et al.* (1995) concluded that toxicity in the San Gabriel River watershed was likely due to non-polar organics and possibly ammonia. This is not unexpected as there are multiple WRPs discharging to the San Gabriel River; these treated effluent discharges comprise roughly 80% of flow during the dry season, contributing as much as 99% of the total

ammonia input (Ackerman *et al.* 2005). In 1992/93, ammonia levels averaged over 10 mg/L. In 2003, however, the WRPs fully implemented nitrification and denitrification treatment (NDN) processes, which subsequently reduced discharged ammonia levels more than 80% to an average of less than two mg/L (Figure 5). Thus, a reduction in toxicity for reaches in the San Gabriel River watershed dominated by WRP effluents can be easily explained.

The lack of toxicity observed in the current study is consistent with other toxicity data collected in recent years. In 2005, a probability-based watershed survey was conducted in the entire San Gabriel River watershed, and 7% of the stream-miles were considered toxic to *Ceriodaphnia* (Stein and Bernstein, unpublished data). Even this toxicity, however, was eliminated after a TIE and subsequent follow-up investigations helped identify and eliminate the illicit discharge responsible.

A second example of reduced toxicity in recent years was observed in routine toxicity monitoring required in the vicinity of the WRPs as a part of their National Pollutant Discharge Elimination System (NPDES) permit requirements. Between June 2003 and June 2006, only 14% of the 269 total samples from 14 different sites exhibited toxicity (i.e., greater than 25% response relative to controls). For this period, toxicity was largely constrained to Coyote Creek (6% of total number of samples) and the uppermost portions of San Jose Creek (6% of total number of samples). Coyote Creek is the same trib-



**Figure 5.** Time series plot of ammonia concentrations in final effluent and receiving water immediately downstream of the Los Coyotes WRP in the lower San Gabriel River Watershed. NDN plant upgrades were completed in June 2003.



utary in which the current study found intermittent toxicity. The uppermost section of San Jose Creek was not monitored during the current study.

In contrast to the main stem of the San Gabriel River where significant resources have been expended to reduce pollutant inputs and minimize toxicity, much less effort has been spent on identifying and remediating sources of toxic pollutants in the Coyote Creek subwatershed. As a result, the toxicity in Coyote Creek has remained. The frequency of toxicity in Coyote Creek has remained similar between 1992/93 and 2004/05; roughly 12 to 22% of the samples were considered toxic. Pesticides available for application by homeowners continue to be one toxicant of concern. Diazinon was identified in 2004/05 (this study), as well as in the 1992/93 study (Bailey *et al.* 1995). The toxicity observed in urban runoff-dominated reaches during this study was intermittent, which is consistent with contributions by homeowner pesticide use (Schiff and Tiefenthaler 2003), illegal/illicit discharges, and observations in other dry weather runoff toxicity studies (Greenstein *et al.* 2004).

## LITERATURE CITED

- Ackerman, D., E.D. Stein and K.C. Schiff. 2005. Dry-season water quality in the San Gabriel River watershed. *Bulletin of the Southern California Academy of Sciences* 104:125-145.
- Bailey, H., A. Villalobos, E. Gottl, L. Brattin, D. Hanes, and D. Hinton. 1995. Site specific study for effluent dominated streams (San Gabriel River, Santa Clara River, Calleguas Creek). Prepared for the California Regional Water Quality Control Board, Los Angeles Region. University of California, Davis. Davis, CA.
- Bay, S.M., D.J. Greenstein, S.-L. Lau, M.K. Stenstrom and C.G. Kelley. 1996. Toxicity of dry weather flow from the Santa Monica Bay Watershed. *Bulletin of Southern California Academy of Sciences* 95:33-45.
- Brown, J. and S. Bay. 2003. Organophosphorus pesticides in the Malibu Creek watershed. Technical Report 403. Southern California Coastal Water Research Project. Westminster, CA.
- Greenstein, D.J., L. Tiefenthaler and S. Bay. 2004. Toxicity of parking lot runoff after application of simulated rainfall. *Environmental Contamination and Toxicology* 47:199-206.
- Los Angeles County Department of Public Works (LACDPW). 2005. Los Angeles County 1994-2005 integrated receiving water impact report. Los Angeles County Department of Public Works. Alhambra, CA.
- Norberg-King, T.J., L.W. Ausley, D.T. Burton, W.L. Goodfellow, J.L. Miller and W.T. Waller. 2005. Toxicity reduction and toxicity identification evaluations for effluents, ambient waters, and other aqueous media. Society of Environmental Toxicology and Chemistry (SETAC) Press. Pensacola, FL.
- Schiff, K. and L. Tiefenthaler. 2003. Contributions of organophosphorus pesticides from residential land uses during dry and wet weather. Technical Report 406. Southern California Coastal Water Research Project. Westminster, CA.
- Stein, E. and L.L. Tiefenthaler. 2004. Characterization of dry weather metals and bacteria in Ballona Creek. Technical Report 427. Southern California Coastal Water Research Project. Westminster, CA.
- United States Environmental Protection Agency (USEPA). 1991. Methods for aquatic toxicity identification evaluation: Phase I toxicity characterization procedures, Second Edition. EPA 600/6-91/003. USEPA, Environmental Research Laboratory. Duluth, MN.
- USEPA. 1993a. Methods for measuring acute toxicity of effluents and receiving waters to freshwater and marine organisms, Fourth Edition. EPA 600/4-90/027. USEPA, Environmental Research Laboratory. Duluth, MN.
- USEPA. 1993b. Methods for aquatic toxicity identification evaluation: Phase II toxicity identification procedures for acute and chronic toxicity, Second Edition. EPA 600/R-92/080. USEPA, Environmental Research Laboratory. Duluth, MN.

## ACKNOWLEDGMENTS

Toxicity testing was conducted by the Los Angeles County Sanitation Districts (Jay P. Bottomley and the staff of the San Jose Creek Water Quality Laboratory) and Nautilus Environmental, Inc. (Chris Stransky, Howard Bailey and staff). Wet weather sampling was conducted by MACTEC Engineering, Inc. (Jay Shrake and staff). Dry weather

sampling was conducted by the Los Angeles County Sanitation Districts (Misty Rogalski and Jeaneal Davis) and the Southern California Coastal Water Research Project (Dario Diehl, David Tsukada, Dawn Petschauer and Diana Young). Information management was provided by Southern California Coastal Water Research Project (Jeff Brown and Larry Cooper). This project was partially supported by the Los Angeles Regional Water Quality Control Board.