

SCCWRP Annual Report 2009

Rapid QPCR-based assay for fecal *Bacteroides* spp. as a tool for assessing fecal contamination in recreational waters

Reagan R. Converse¹, A. Denene Blackwood¹, Marek Kirs², Rachel T. Noble¹ and John F. Griffith

¹University of North Carolina at Chapel Hill, Institute of Marine Sciences, Morehead City, NC

²Cawthron Institute, Nelson, New Zealand

ABSTRACT

Concentrations of fecal indicator bacteria (e.g., *Escherichia coli* and *Enterococcus* sp.) can only be used in limited ways for determining the source of fecal contamination in recreational waters because they cannot distinguish human from non-human fecal contamination. Several *Bacteroides* spp. have been suggested as potential alternative indicators. We have developed a rapid, culture-independent method for quantifying fecal *Bacteroides* spp. using quantitative polymerase chain reaction (QPCR) targeting the 16S rRNA gene. The assay specifically targets and quantifies the most common human *Bacteroides* spp. The details of the method are presented, including analyses of a wide range of fecal samples from different organisms. Specificity and performance of the QPCR assay were also tested via a laboratory experiment in which human sewage and gull guano were inoculated into a range of environmental water samples. Concentrations of fecal *Bacteroides* spp., total *Enterococcus* sp., *Enterococcus faecium*, *Enterococcus faecalis*, and *Enterococcus casseliflavus* were measured using QPCR, and total *Enterococcus* sp. and *E. coli* were quantified by membrane filtration (MF). Samples spiked with gull guano were highly concentrated with total *Enterococcus* sp., *E. coli*, *E. faecalis*, and *E. casseliflavus*, demonstrating that these indicators are prominent in animal feces. On the other hand, fecal *Bacteroides* spp. concentrations were high in samples containing sewage and were relatively low in samples spiked with gull guano. Sensitivity and specificity results suggest that the rapid fecal *Bacteroides* spp. QPCR assay may be a useful tool to effectively predict the presence and concentration of human-specific fecal pollution.

Full Text

ftp://ftp.sccwrp.org/pub/download/DOCUMENTS/AnnualReports/2009AnnualReport/AR09_235_247.pdf