SCCWRP Annual Report 2010

A real-time qPCR assay for the detection of the *nifH* gene of *Methanobrevibater smithii*, a potential indicator of sewage pollution

Christopher Johnston, Jennifer A. Unfar, John F. Griffith, Janet A. Gooch and Jill R. Stewart

ABSTRACT

Methanobrevibacter smithii is a methanogenic archaea prevalent in the intestinal tract of humans. Due to its abundance in the human gut and low likelihood of regrowth in the environment, the nifH gene of M. smithii may be a useful marker of sewage pollution in water. This paper describes the development of a real-time quantitative polymerase chain reaction (RT-qPCR) assay to detect the nifH gene of M. smithii. Quantification standards were prepared from M. smithii genomic DNA dilutions and a standard curve used to quantify the target gene and calculate estimated genome equivalency units (GEUs). A competitive internal positive control was designed and incorporated into the assay to assess inhibition in environmental extracts. Testing the assay against a panel of 23 closely related methanogen species demonstrated specificity of the assay for M. smithii. A set of 36 blind water samples was then used as a field test of the assay. The internal control identified varying levels of inhibition in 29 of 36 (81%) samples, and the M. smithii target was detected in all water samples with known sewage input. Our results suggest that the quantitative polymerase chain reaction (qPCR) assay targeting the M. smithii nifH gene developed in this study is both sensitive and rapid, and shows promise as a reliable indicator of sewage in environmental waters.

Full Text

ftp://ftp.sccwrp.org/pub/download/DOCUMENTS/AnnualReports/2010AnnualReport/ar10_127_138.pdf