

## SCCWRP Annual Report 2012

### Passive sampling to measure background dissolved persistent organic pollutant concentrations in the water column of the Palos Verdes Shelf Superfund site

Loretta A. Fernandez<sup>1</sup>, Wenjian Lao, Keith A. Maruya, Carmen White<sup>2</sup> and Robert M. Burgess<sup>1</sup>

<sup>1</sup>United States Environmental Protection Agency, Atlantic Ecology Division, Narragansett, RI

<sup>2</sup>United States Environmental Protection Agency, Region 9, San Francisco, CA

#### ABSTRACT

Passive sampling was used to deduce water concentrations of persistent organic pollutants (POPs) in the vicinity of a marine Superfund site on the Palos Verdes Shelf, California, USA. Pre-calibrated solid phase microextraction (SPME) fibers and polyethylene (PE) strips that were pre-loaded with performance reference compounds (PRCs) were co-deployed for 32 days along an 11 station gradient at bottom, surface and mid-water depths. Retrieved samplers were analyzed for DDT congeners and their transformation products (DDE, DDD, DDMU, and DDNU) and 43 PCB congeners using GC-EI- and NCI-MS. PRCs were used to calculate compoundspecific fractional equilibration achieved *in situ* for the PE samplers, using both an exponential approach to equilibrium model (EAE) and numerical integration of Fickian diffusion (NI) model. The highest observed concentrations were for p,p'-DDE, 2200 pg/L and 990 pg/L deduced from PE and SPME, respectively. The difference in these estimates could be largely attributed to uncertainty in equilibrium partition coefficients, unaccounted for disequilibrium between samplers and water, or different time scales over which the samplers average. The concordance between PE and SPME estimated concentrations for DDE was high ( $R^2 = 0.95$ ). PCBs were only detected in PE samplers, due to their much larger size. Nearbottom waters adjacent to and down current from sediments with the highest bulk concentrations exhibited aqueous concentrations of DDTs and PCBs that exceeded Ambient Water Quality Criteria (AWQC) for human and aquatic health, indicating the need for future monitoring to determine the effectiveness of remedial activities taken against the most highly contaminated surface sediments at this site.

#### Full Text

[ftp://ftp.sccwrp.org/pub/download/DOCUMENTS/AnnualReports/2012AnnualReport/ar12\\_049\\_060.pdf](ftp://ftp.sccwrp.org/pub/download/DOCUMENTS/AnnualReports/2012AnnualReport/ar12_049_060.pdf)