

SCCWRP Annual Report 2012

Nontargeted comprehensive twodimensional gas chromatography/timeof- flight mass spectrometry method and software for inventorying persistent and bioaccumulative contaminants in marine environments

Eunha Hoh¹, Nathan G. Dodder, Steven J. Lehotay², Kristin C. Pangallo³, Christopher M. Reddy⁴ and Keith A. Maruya

¹San Diego State University, Graduate School of Public Health, San Diego, CA

²United States Department of Agriculture, Agricultural Research Service, Eastern Regional Research Center, Wyndmoor, PA

³Colgate University, Department of Chemistry, New York, NY

⁴Woods Hole Oceanographic Institution, Department of Marine Chemistry and Geochemistry, Woods Hole, MA

ABSTRACT

Analytical methods for contaminant monitoring are generally targeted; i.e., they measure defined lists of compounds. Routine monitoring projects using targeted methods are not usually designed to screen for unrecognized or novel contaminants, and therefore miss compounds within the region or population of study that cause, or have the potential to cause, adverse biological impacts. We describe a non-targeted analytical method utilizing direct sample introduction (DSI) coupled to comprehensive two-dimensional gas chromatography with time-offlight mass spectrometry (GC×GC/TOF-MS). To test the capabilities of this instrumental method within the context of marine contaminant surveys, we characterized a broad array of non-polar, persistent and bioaccumulative contaminants in Atlantic common dolphin (*Delphinus delphis*) blubber, including compounds that are not typically monitored. Compound identifications were made by searching a standard reference database, contemporaneously analyzing mass spectra from reference standards, and *de novo* interpretation. We identified a total of 271 compounds belonging to 24 classes; all compounds but one were halogenated. Anthropogenic contaminants and halogenated natural products were concurrently detected. Eighty-six compounds were anthropogenic contaminants that are not routinely targeted in environmental surveys, and 54 compounds were halogenated natural products. One hundred twelve spectra were identified *de novo*, demonstrating that exclusive reliance on commercially available reference standards and mass spectral libraries may miss a significant fraction of identifiable compounds. We also cataloged 27 halogenated mass spectra that were not able to be identified. Due to the volume and complexity of the identification data, we developed custom software to organize and provide shared access to the identified mass spectra and related information. The non-targeted analytical method and data reporting system, in combination with the analysis of a high-trophic level sentinel species, demonstrates a framework for creating an inventory of persistent and bioaccumulative contaminants in marine environments, with the future goal of suggesting new compounds for further investigation by targeted monitoring and risk assessment.

Full Text

ftp://ftp.sccwrp.org/pub/download/DOCUMENTS/AnnualReports/2012AnnualReport/ar12_073_084.pdf