SCCWRP Annual Report 2012

Reach-scale geomorphic and biological effects of localized stream bank armoring

Eric D. Stein, Matthew R. Cover¹, A. Elizabeth Fetscher, Clare O'Reilly², Roxana Guardado1 and Christopher W. Solek

¹California State University Stanislaus, Turlock, CA

ABSTRACT

Armoring of stream banks is a common management response to perceived threats to adjacent infrastructure from flooding or erosion. Despite their pervasiveness, the effects of reach-scale bank armoring have received less attention than those of channelization or watershed-scale hydromodification. In this study we explored mechanistic ecosystem responses to armoring by comparing conditions upstream, within, and downstream of six stream reaches with bank armoring in Southern California. Assessments were based on four common stream-channel assessment methods: 1) traditional geomorphic measures, 2) the California Rapid Assessment Method (CRAM) for wetlands, 3) bioassessment with benthic macroinvertebrates, and 4) bioassessment with stream algae. Although physical responses varied among stream types (mountain, transitional, and lowland), armored segments generally had lower slopes, more and deeper pools and fewer riffles, and increased sediment deposition. Several armored segments exhibited channel incision and bank toe failure. All classes of biological indicators showed subtle, mechanistic responses to physical changes. However, extreme heterogeneity among sites, the presence of catchment-scale disturbances, and low sample size made it difficult to ascribe observed patterns solely to channel armoring. The data suggest that species-level or functional group-level metrics may be more sensitive tools than integrative indices of biotic integrity (IBIs) to local-scale effects.

Full Text

ftp://ftp.sccwrp.org/pub/download/DOCUMENTS/AnnualReports/2012AnnualReport/ar12 113 128.pdf

²California Ocean Protection Council, Oakland, CA