

SCCWRP Annual Report 2012

Thresholds of adverse effects of macroalgal abundance and sediment organic matter on benthic habitat quality in estuarine intertidal flats

Martha Sutula, Lauri Green¹, Giancarlo Cicchetti², Naomi Detenbeck² and Peggy Fong¹

¹University of California, Department of Evolutionary Biology and Ecology, Los Angeles, CA

²US Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Atlantic Ecology Division, Narragansett, RI

ABSTRACT

Confidence in the use of macroalgae as an indicator of estuarine eutrophication is limited by the lack of quantitative data on the thresholds of adverse effects of macroalgae on benthic habitat quality. In the present study, we utilized sediment profile imagery (SPI) to identify thresholds of adverse effect of macroalgal biomass, sediment organic carbon (%OC) and sediment nitrogen (%N) concentrations on the apparent Redox Potential Discontinuity (aRPD), the depth that marks the boundary between oxic near-surface sediment and the underlying suboxic or anoxic sediment. We surveyed 16 sites in eight California estuaries. At each site, SPI, macroalgal biomass and sediment cores were collected at 20 locations along an intertidal transect; cores were analyzed for %OC and %N. Classification and Regression Tree (CART) analysis was used to identify step thresholds associated with a transition from “reference” or natural background levels of macroalgae, defined as that range in which no effect on aRPD was detected. Ranges of 3 to 15 g dw macroalgae m⁻², 0.4 to 0.7% OC and 0.05 to 0.07% N were identified as transition zones from reference conditions across these estuaries. Piecewise regression analysis was used to identify thresholds of adverse effects, associated with the transition from a steep decline in aRPD to a consistent minimum value. Levels of 175 g dw macroalgae m⁻², 1.1% OC and 0.1% N were identified as thresholds of adverse effects, associated with a shallowing of aRPD to near zero depths. As an indicator of ecosystem condition, shallow aRPD has been related to reduced volume and quality for benthic infauna and alteration in community structure. These effects have been linked to reduced availability of forage for fish, birds and other invertebrates, as well as to undesirable changes in sediment biogeochemical processing of nutrients.

Full Text

ftp://ftp.sccwrp.org/pub/download/DOCUMENTS/AnnualReports/2012AnnualReport/ar12_189_210.pdf