



APPENDIX F:


Regressions of Community Characteristics
Versus Latitude and Depth




Appendix F1. Community characteristics of the Southern California Bight vs. latitude:
a) number of taxa, **b)** Shannon-Wiener diversity index, **c)** dominance, **d)** evenness,
e) total abundance / m^2 , and **f)** total biomass / m^2 . * = significant.




Appendix F2. Proportion of infaunal abundance / sample of six taxonomic groups on the Southern California Bight vs. latitude: a) annelida, b) arthropoda, c) ophiuroidea, d) misc. echinodermata, e) mollusca and f) other phyla. * = significant.




Appendix F3. Proportion of infaunal biomass (gms wet weight) / sample of six taxonomic groups on the Southern California Bight vs. latitude: a) annelida, b) arthropoda, c) ophiuroidea, d) misc. echinodermata, e) mollusca and f) other phyla.
 $*$ = significant.



Appendix F4. Community characteristics of the Southern California Bight vs. depth:
a) number of taxa, **b)** Shannon-Wiener diversity index, **c)** dominance, **d)** evenness,
e) total abundance / m², and **f)** total biomass / m². * = significant.



Appendix F5. Proportion of infaunal abundance / sample of six taxonomic groups on the Southern California Bight vs. depth: a) annelida, b) arthropoda, c) ophiuroidea, d) misc. echinodermata, e) mollusca and f) other phyla. * = significant.



Appendix F6. Proportion of infaunal biomass (gms wet weight) / sample of six taxonomic groups on the Southern California Bight vs. depth: a) annelida, b) arthropoda, c) ophiuroidea, d) misc. echinodermata, e) mollusca, and f) other phyla.
* = significant.