

APPENDIX I

Coastal water quality impact of storm water runoff from an urban watershed in southern California

Jong Ho Ahn¹, Stanley B. Grant^{1,*}, Cristiane Q. Surbeck¹, Paul M. DiGiacomo², Nikolay P. Nezlin³, Sunny Jiang⁴

¹Department of Chemical Engineering and Materials Science
Henry Samueli School of Engineering, University of California, Irvine, CA 92697
²Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109

³Southern California Coastal Water Research Project, Westminster, CA 92683

⁴Department of Environmental Health, Science, and Policy

School of Social Ecology, University of California, Irvine, CA 92697

Abstract

Field studies were conducted to assess the coastal water quality impact of storm water runoff from the Santa Ana River, which drains a large urban watershed located in southern California. Storm water runoff from the river leads to very poor surf zone water quality, with fecal indicator bacteria concentrations exceeding California ocean bathing water standards by up to 500 %. However, cross-shore currents (e.g., rip cells) dilute contaminated surf zone water with cleaner water from offshore, such that surf zone contamination is generally confined to < 5 km around the river outlet. Offshore of the surf zone, storm water runoff ejected from the mouth of the river spreads out over a very large area, in some cases exceeding 100 km² based on satellite observations. Fecal indicator bacteria concentrations in these large storm water plumes generally do not exceed California ocean bathing water standards, even in cases where offshore samples test positive for human pathogenic viruses (human adenoviruses and enteroviruses) and fecal indicator viruses (F⁺ coliphage). Multiple lines of evidence indicate that bacteria and viruses in the offshore storm water plumes are either associated with relatively small particles (< 53 µm) or not particle-associated. Collectively, these results demonstrate that storm water runoff from the Santa Ana River negatively impacts coastal water quality, both in the surf zone and offshore. However, the extent of this impact, and its human health significance, is influenced by numerous factors, including prevailing ocean currents, within-plume processing of particles and pathogens, and the timing, magnitude and nature of runoff discharged from river outlets over the course of a storm.

Introduction

Oceans adjacent to large urban areas, or "urban oceans", are the final repositories of pollutants from a myriad of point and non-point sources of human waste (1). Pollutants are transported to the urban ocean by surface water runoff (1-4), discharge of treated sewage through submarine outfalls (5), wet and dry deposition of airborne pollutants (6), and submarine discharge of contaminated groundwater (7). Until recently, effluent from sewage treatment plants was often the primary source of urban coastal pollution -- including nutrients, pathogens, pesticides, and heavy metals (8). However, pollutant loading from many sewage treatment plants has declined over the past several decades due to improvements in civil infrastructure (e.g., separation of the storm and sanitary sewer systems to prevent combined sewer overflows), pollutant source control, and disposal/treatment technology (9). As a result, surface water runoff, in many cases, has supplanted sewage treatment plants as the primary source of pollutant loading to the urban ocean (3, 10).

The focus of this study is the coastal water quality impact of surface water runoff during storms -- or "storm water runoff" -- from an urban watershed in southern California. The study was motivated by several considerations. First, beneficial use designations for the coastal ocean in southern California apply year-round and, consequently, watershed managers are legally required to develop storm water management plans for reducing wet-weather impairments of the coastal ocean (11). The impact of storm water runoff on coastal water quality is of particular concern in arid regions like southern California because, on an annual basis, a large percentage (> 99.9% according to Reeves et al. (2) and > 95% according to Schiff et al. (10)) of the surface water runoff and associated pollution flows into the ocean during a few storms in the winter. Second, while recreational use of the coastal ocean in southern California is lighter in the winter, compared to the summer, winter ocean recreation is still very common, particularly among surfers who surf the large waves that often accompany storm events (R. Wilson, personal communication). Third, to the extent that particles in storm water runoff are associated with pathogens and other contaminants, their discharge to the ocean during storms may serve as a source of near shore pollution that persists long after the storm season is over (10, 12). Finally, in many urban watersheds in southern California and elsewhere, the flow of storm water runoff in urban watersheds is highly regulated by civil infrastructure (e.g., dams) designed to minimize flood potential and maximize water reclamation. As will be demonstrated later in this paper, the regulated nature of storm water runoff implies that the ocean discharge of storm water runoff from urban watersheds can occur days after the cessation of rain, when the potential for human exposure to pathogens by marine recreational contact is significant.

This paper describes how storm water runoff from several major rivers in southern California - with particular focus on the Santa Ana River in Orange County -- impacts coastal water quality, as measured by turbidity, particle size spectra, total organic carbon, fecal indicator bacteria, fecal indicator viruses, and human pathogenic viruses. The present study is unique in the combination of data resources utilized, including data and information from routine surf zone water quality and wave field monitoring programs, an automated in-situ ocean observing sensor, shipboard sampling cruises, and satellite sensors. Further, this is the first wet weather study to examine the linkage between water quality in the surf zone -- where routine monitoring samples are collected and most human exposure occurs -- and water quality offshore of the surf zone. The work described in this study was carried out in parallel with a watershed-focused study that examined the spatial variability of fecal indicators, and the relationship between suspended particle size

and fecal indicators, in storm runoff from the Santa Ana River watershed (13). Background information is available elsewhere on coastal water quality impairment at our Orange County field site (2, 14-18) and the transport and mixing dynamics of sediment plumes as they flow into the coastal ocean from river outlets in southern California (4, 19, 20).

Materials and Methods

Rainfall and river discharge. Weather information and Next Generation Radar (NEXRAD) images for planning the field studies and interpreting rainfall patterns, were obtained online from the National Weather Service (<http://www.nwsla.noaa.gov/>). Precipitation and stream discharge data were obtained at two sites, one located where the Santa Ana River crosses 5th Street in the City of Santa Ana, and another located where the San Gabriel River crosses Spring Street in the City of Long Beach (black squares in inset, Figure 1). These data were obtained, respectively, from the U.S. Army Corps of Engineers and the Los Angeles County Department of Public Works. Both of these gauge sites are located relatively close (within 11 km) to the rivers' respective ocean outlets, and hence stream flow measured at these sites will likely make its way to the ocean.

Surf zone measurements: NEOCO data. Time series of water temperature, conductivity, chlorophyll, and water depth were obtained from an instrument package deployed at the end of the Newport Pier, where the local water depth is between 6.5 and 9 m (blue star in Figure 1). This instrument package is part of a recently deployed network of coastal sensors in southern California called the Network for Environmental Observations of the Coastal Ocean (NEOCO). The NEOCO sensor package contains a SBE-16plus CTD (Sea-Bird Electronics, Inc., Bellevue, WA) and a Seapoint Chlorophyll Fluorometer (Seapoint Sensors, Inc.). These instruments are mounted on a pier piling at a depth of approximately 1m (below mean lower low water) and programmed to acquire data at a sampling frequency of 0.25 min⁻¹.

Surf zone measurements: fecal indicator bacteria and breaking waves. The concentration of fecal indicator bacteria in the surf zone was measured at 17 stations (black circles along shoreline in Figure 1) by personnel at the Orange County Sanitation District (OCSD). The stations are designated by OCSD according to their distance (in thousands of feet) north or south of the Santa Ana River outlet (e.g., station 15N is located approximately 15,000 ft, approximately 5 km, north of the Santa Ana River outlet). Water samples were collected five days per week (not on Friday and Sunday) from 5:30 to 10:00 local time at ankle-depth on an incoming wave, placed on ice in the dark, and returned to the OCSD (Fountain Valley, CA) where they were analyzed within 6 h of collection for total coliform (TC), fecal coliform (FC), and enterococci bacteria (ENT) using standard methods 9221B and 9221E, and EPA method 1600, respectively. Results are reported in units of colony forming units per 100 mL of sample (CFU/100mL). Wave conditions, including both the direction and height of breaking waves, were recorded by lifeguards at the Newport Beach pier (near surf zone station 15S, Figure 1) twice per day, once at 7:00 and again at 14:00 local time.

Offshore measurements: satellite ocean color imagery. The satellite images used in this study were collected by NASA's Moderate-Resolution Imaging Spectroradiometer (MODIS) instruments. These instruments operate onboard two near-polar sun-synchronous satellite platforms orbiting at 705 km altitude: Terra (since 24 February 2000) and Aqua (since 24 June

2002). Terra passes across the equator from north to south at ~ 10:30 local time, while Aqua passes the equator south to north at ~ 13:30 local time. As such, all the images were acquired within two hours before or after local noon, or between 18:00 and 22:00 UTC. The MODIS sensors collect data in 36 spectral bands, from 400 to 14,000 nm. We utilized bands 1 (250 m spatial resolution, 620 - 670 nm), 3, and 4 (500 m resolution, 459 - 479 and 545 - 565 nm, respectively) to produce “true color” (i.e., RGB) images, with band 1 used for the Red channel, band 4 for the Green channel, and band 3 for the Blue channel. Using a MATLAB program, the 500 m Green (band 4) and Blue (band 3) monochrome channels were “sharpened” to 250 m resolution using fine details from the higher resolution Red channel (band 1). Then, the contrast of each of these monochrome channels was increased to emphasize maximum details in the coastal ocean region of interest. Finally, all three monochrome channels (i.e. Red, Green, and Blue) were combined to form a single true color image. In all, sixteen satellite images from February 23 to March 5 were acquired and processed for this study; four of them were selected as most illustrative, based on their quality and observed features. The timing of these satellite acquisitions relative to the storms and sampling periods is indicated at the top of Figure 2.

Offshore measurements: sampling cruises. The offshore monitoring grid (red triangles in Figure 1) was sampled during three separate cruises on 23 February, 28 February, and 1 March 2004, coinciding with a sequence of storm events in late February 2004. Table 1 provides a summary of activities performed during each cruise. A short description of the offshore sampling and analysis protocols is presented here; details can be found in the Supporting Information for this paper. All offshore water samples were analyzed for salinity and fecal indicator bacteria -- specifically, total coliform (TC), *Escherichia coli* (EC, a subset of FC), and enterococci bacteria (ENT) -- using the defined substrate tests known commercially as Colilert and Enterolert implemented in a 97-well quantitray format; results are reported in units of most probable number of bacteria per 100 mL of sample (MPN/100 mL). A subset of the offshore water samples was analyzed for total organic carbon (TOC) by USEPA Method 415.1, fecal indicator viruses (F^+ coliphage) by a two-step enrichment method (USEPA Method 1601), and human pathogenic viruses (human adenovirus and human enterovirus) by real-time quantitative polymerase chain reaction (Q-PCR), nested PCR, and reverse-transcriptase (RT)-PCR using published protocols (21-25). Details on the PCR protocols used here can be found in the Supporting Information for this paper.

Coincident with the collection of the offshore water samples, temperature, particle size spectra, and light transmissivity were measured using a LISST-100 (Laser In-Situ Scattering and Transmissometry) analyzer (Sequoia Scientific, In., Bellevue, WA). The LISST-100 estimates the particle volume per unit fluid volume (ΔV) resident in 32 logarithmically-spaced particle diameter bins ranging in size from $d_p = 2.5$ to 500 μm . At least ten replicates of the particle size spectra were collected at each offshore station. Following the recommendation of Mikkelsen (26), ΔV was taken as the median of all replicate measurements. The LISST-100 data are presented in this paper in one of three ways: (1) particle size spectra represented by plots of $\Delta V / \Delta \log d_p$ against $\log d_p$, (2) the number of particles per unit fluid volume or total number concentration (TNC), and (3) the number-averaged particle size, \bar{d} . The last two parameters were computed from the particle size spectra as follows (26, 27):

$$TNC = \sum_{i=1}^{32} \frac{6\Delta V_i}{\pi d_{p,i}^3}$$

(1a)

$$\bar{d} = \sqrt[3]{\frac{6}{\pi} \frac{\sum_{i=1}^{32} \Delta V_i}{TNC}}$$

(1b)




Figure 1. Map showing location of field site and sampling sites in the surf zone and offshore. Also shown are the locations of the NEOCO sensor on the end of the Newport Pier, and rain and stream gauges located on the Santa Ana River and San Gabriel River. Abbreviations are Los Angeles River (LAR), San Gabriel River (SGR), Santa Ana River (SAR), Orange County Sanitary District (OCSD), and University of California, Irvine (UCI).

Table 1. Summary of analyses performed during the sampling cruises.

Sampling Parameters	Methods	Number of Offshore Sites Sampled			
		February 23 2004	February 28 2004	March 1 2004	
Conductivity¹	Thermo Orion162A or CTD (SBE-32)	20	21	21	
Temperature²	Thermocouple w/ LISST-100 or CTD (SBE-32)	20	21	21	
Total coliform, <i>Escherichia coli</i>, Enterococcus³	Colilert and Enterolert (IDEXX)	20 (+ 2 sets of fractionated samples)	21 (+ 6 sets of fractionated samples)	21	
Total Organic Carbon⁴	EPA 415.1	17 (+ 2 sets of fractionated samples)	-	-	
Human Adenoviruses & Enteroviruses⁵	Nested PCR RT-PCR	2	6	-	
Fecal Indicator Viruses (F⁺coliphage)⁵	Two-step Enrichment	2	6	-	
Particle Size Spectra	LISST-100 (Light Diffraction)	20	16	21	
Transmissivity	LISST-100	20	16	21	

¹ Measured using a Thermo Orion 162A conductivity meter on 23 February; a CTD instrument (SBE-32) on 28 February and 1 March.

² Measured using a thermocouple bundled with a LISST-100 on 23 February; a CTD instrument (SBE-32) on 28 February and 1 March.

³ Samples collected by UCI and analyzed by OCSD on 23 February; collected and analyzed by OCSD on 28 February and 1 March. Fractionated samples collected and analyzed by UCI on 23 and 28 February.

⁴ Collected by UCI and analyzed by Del Mar Analytical.

⁵ Carried out on the fractionated samples, and measured using a real time PCR for enterovirus and a nested PCR for adenovirus, respectively.

Results and Discussion

Rainfall and river discharge. Over the period of study (18 February through 3 March 2004), four rain events were recorded by the rain gauge on the Santa Ana River in the City of Santa Ana (black curve, top panel, top axis, Figure 2). The first event accumulated 16.0 mm of rain in the afternoon of 21 February (RE₁ in Figure 2), the second event accumulated 23.4 mm of rain in the afternoon of 22 February (RE₂), the third event accumulated 51.3 mm of rain in the evening of 25 February (RE₃), and the fourth event accumulated 6.8 mm of rain in the evening of 1 March (RE₄). The rain gauge located on the San Gabriel River in the City of Long Beach did not record RE₂, and recorded a fifth rain event on 18 February (red curve, top panel, top axis, Figure 2). The difference in rainfall recorded at the Santa Ana River and San Gabriel River sites is a consequence of the spatial variability of rainfall near the coast (see Figures S1-S2, Supporting Information, for NEXRAD maps acquired during RE₁ and RE₂). Records of stream discharge (in units of m³/s) at the Santa Ana River and San Gabriel River sites are also quite different (black and red curves, top panel, bottom axis, Figure 2). While rainfall and stream discharge are coupled at the San Gabriel River site (i.e., stream discharge increases shortly after locally recorded rain events, compare set of red curves in top panel, Figure 2), rainfall and stream discharge are frequently uncoupled at the Santa Ana River site. For example, the Santa Ana River discharge events DE₃ and DE₄ do not obviously correlate with records of local rainfall. Instead, these two discharge events can be traced to storm water runoff generated from inland regions of the Santa Ana River watershed that was released from inland dams after the cessation of rain (13). For comparison we have also included in the plot hourly volume discharge records (unit of m³/s, blue curve, top panel, Figure 2) of treated sewage discharged from the Orange County Sanitation District (OCSD) sewage outfall (courtesy of OCSD).

Surf zone measurements: NEOCO data. Water level, salinity, temperature, and chlorophyll measurements at the NEOCO sensor -- located on the end of the Newport Pier at the offshore edge of the surf zone -- are presented in Figure 2 (second and third panels). The largest rain event (RE₃) and the largest discharge of storm water runoff from the Santa Ana River (DE₄) occurred during a neap tide when the daily tide range was small (see quarter moon and tide level measurements in the second panel, Figure 2). The other rainfall and stream discharge events occurred during periods of time when the daily tide range was larger, either during the transition from spring to neap tide (RE₁, RE₂, DE₁, DE₂, DE₃), or during the transition from neap to spring tide (RE₄, DE₅).

Salinity recorded at the NEOCO sensor is characterized by a series of low salinity events, relative to ambient ocean water salinity of 32.5 to 33.0 ppt (salinity events SE₁ through SE₆, Figure 2). These low salinity events may be caused, at least in part, by storm water discharged from the Santa Ana River (e.g., SE₆ appears to be related to DE₄). However, correlating discharge and the low salinity events is complicated by the fact that once river water is discharged to the ocean, its offshore transport is controlled by a complex set of near shore currents (28). These near shore currents, and their impact on the spatial distribution of storm water runoff plumes, are explored in the next several sections. Temperature and chlorophyll records at the NEOCO sensor appear to be relatively unaffected by rainfall and/or discharge from the Santa Ana River. Surf zone temperature exhibits a diurnal pattern consistent with solar heating (i.e., temperatures are higher during the day and lower at night). Chlorophyll measurements indicate a bloom event occurred early in the study period (Bloom Event 1, BE₁), but this bloom event mostly dissipated prior to the rain and discharge events that occurred later.

While the chlorophyll fluorometer was being maintained during this period, we cannot rule out the possibility that the downward trend in the chlorophyll signal is related to instrument fouling.




Figure 2. Timing of the satellite images (blue lettering) is shown at the top and black squares indicate offshore sampling cruises. Time series measurements of rainfall, stream discharge at the Santa Ana River and San Gabriel River, and discharge of treated sewage from the OCSD outfall (top panel); water level, salinity, temperature, and chlorophyll measured at the NEOCO sensor (second and third panels); the direction and height of breaking waves at the Newport Beach Pier (fourth panel); and the concentration of fecal indicator bacteria in the surf zone (color contour plots, fifth through seventh panels).

Surf zone measurements: wave data and along-shore currents. Wave conditions, including the direction and height of breaking waves, were recorded twice per day by lifeguards stationed at the Newport Pier (surf zone station 15S, Figure 1). These wave data, which are plotted in the fourth panel of Figure 2, can be divided into five events, depending on whether waves approach the beach from the west (WE₁, WE₃, and WE₅) or from the south to southwest (WE₂ and WE₄). Because this particular stretch of shoreline strikes northwest-southeast (see Figure 1), waves approaching the beach from the west are likely to yield a down-coast surf zone current (i.e., directed to the southeast). Likewise, waves approaching the beach from the south are likely to yield an up-coast surf zone current (i.e., directed to the northwest) (28, 29).

This expectation is consistent with the salinity signal measured at the NEOCO sensor, which is located approximately 5 km down-coast of the Santa Ana River ocean outlet. The onset of low salinity event SE₆ at the NEOCO sensor coincides very closely in time with the change in wave conditions from WE₂ to WE₃, and a likely change in the direction of the surf zone current from up-coast to down-coast (Figure 2). Discharge from the Santa Ana River was particularly high during this period (note that discharge event DE₄ overlaps wave events WE₂ and WE₃). Hence the onset of SE₆ was probably triggered by a change in the direction of wave-driven surf zone currents from up-coast during WE₂ to down-coast during WE₃ and a consequent down-coast transport of storm water runoff entrained in the surf zone from the Santa Ana River during DE₄.

Employing the same logic, low salinity events SE₃ through SE₅, which occurred during a period when waves were out of the south to southwest, may have originated from storm water discharged by river outlets and/or embayment located down-coast of the NEOCO sensor (e.g., the Newport Bay outlet). Low salinity events SE₁ and SE₂, which occurred during a period when waves were out of the west, may have originated from storm water discharged by outlets located up-coast of the NEOCO sensor, although no significant discharge from the Santa Ana River was recorded during this period of time.

It should be noted that some of these low salinity events may have originated from the cross-shore transport of lower salinity water from offshore -- perhaps from surface runoff plumes, or submarine waste water fields associated with local sewage outfalls (16) -- and/or from the submarine discharge of low salinity ground water (7). While the power-plant cooling water intake and outfall appear to affect local circulation patterns offshore of Huntington Beach (30), the power-plant effluent consists of pure ocean water and therefore is very unlikely to be a source of the low salinity events documented in Figure 2. It is theoretically possible that the OCSD sewage outfall is a source of SE₁ and SE₂, although there is nothing unusual about the sewage discharge rates observed during these two periods of time (compare SE₁ and SE₂ with the blue curve, top panel, Figure 2).

Surf zone measurements: fecal indicator bacteria. The concentrations of the three fecal indicator bacteria groups (TC, FC, and ENT) in the surf zone are presented as a set of color contour plots in Figure 2 (bottom three panels). Fecal indicator bacteria concentrations were log-transformed in order to visualize the temporal and spatial variability associated with these measurements. For comparison, the California single-sample standards for the three fecal indicator bacteria (10^4 for TC, $10^{2.602}$ for FC, and $10^{2.017}$ for ENT, all CFU or MPN/100 mL) are indicated by a set of arrows on the scale bar in the figure. The concentration of fecal indicator bacteria was frequently elevated around the ocean outlet of the Santa Ana River (near surf zone

station 0), particularly during and after rain events when storm water was discharging from the river. For example, during storm water discharge events (DE₃ and DE₄), water quality around the Santa Ana River outlet was very poor (see water quality events TC₂, FC₂, and ENT₂ in Figure 2). During this period of time, fecal indicator bacteria concentrations around the Santa Ana River outlet frequently exceeded one or more State standards, in some cases by as much as 300 to 500 % (depending on the fecal indicator group).

The spatial distribution of fecal indicator bacteria in the surf zone around the Santa Ana River outlet appears to be controlled by local wave conditions, in a manner consistent with the earlier discussion of wave-driven surf zone currents. When waves approach the beach from the west and down-coast currents are likely to prevail, the concentration of fecal indicator bacteria in the surf zone is higher on the down-coast side of the ocean outlet (compare WE₁ with TC₁, FC₁, ENT₁ and WE₃ with TC₃, FC₃, ENT₃). Likewise, when waves approach the beach from the south and up-coast currents are likely to prevail, the concentration of fecal indicator bacteria in the surf zone is higher on the up-coast side of the ocean outlet (compare WE₂ with TC₂, FC₂, ENT₂). The exception is a short period of time when relatively small waves (wave height < 0.5 m) approach the beach from the southwest and the concentration of fecal indicator bacteria is higher on the down-coast side of the river (compare WE₄ with TC₄, FC₄, ENT₄). This exception can be rationalized by noting that waves out of the southwest break with their crests parallel to the beach, and hence the direction of long-shore transport in the surf zone is likely to be unpredictable under these conditions. The apparent time delay between change in wave direction (e.g., from WE₁ to WE₂) and change in the spatial distribution of fecal indicator bacteria around the Santa Ana River outlet (e.g., from TC₁ to TC₂) is, at least in part, a sampling artifact. Wave height and direction were recorded twice per day while fecal indicator bacteria concentrations in the surf zone were sampled at most once per day (the gray dots in the color contour plots indicate the timing of surf samples at each station).

Storm water runoff discharged from the Santa Ana River appears to severely impact water quality in the surf zone over a fairly limited stretch of the beach (< 5 km either side of the river between surf zone stations 15N and 15S). This spatial confinement of storm water plumes in the surf zone, which is particularly evident for FC and ENT, could be the result of physical transport processes (e.g., dilution by rip cell mediated exchange of water between the surf zone and offshore) and/or non-conservative processes (e.g., the removal of fecal indicator bacteria from the surf zone by die-off and/or sedimentation) (28, 29). An analysis of historical fecal indicator bacteria measurements at Huntington Beach concluded that the length of surf zone impacted by point sources of fecal indicator bacteria, such as the Santa Ana River, is influenced more by rip cell dilution, and less by non-conservative processes such as die-off (31). The decay length scale reported here of 5 km is very close to the length scale predicted by rip cell dilution alone (2 - 4 km, assuming a rip cell spacing of 0.5 km) (31). Hence, die-off probably plays a secondary role, compared to dilution, in limiting the distance over which water quality is impaired in the surf zone by storm water runoff from the Santa Ana River.

Fecal indicator bacteria events also occur in the surf zone at the northern (events TC₆, TC₇, ENT₆, ENT₇) and southern (events TC₅, FC₅, and ENT₅) edges of our study area. Possible sources of these fecal indicator bacteria events include storm water discharged from the Huntington Harbor and Newport Bay Harbor located at the extreme northern (5 km up-coast of station 39N) and southern (stations 27S and 29S) ends of the study site and, possibly, from river outlets located outside of the study area (e.g., the Los Angeles River and San Gabriel River, see

inset in Figure 1). Boehm and co-workers (32, 33) suggested that the OCSD sewage outfall might be a source of fecal indicator bacteria in the surf zone at Huntington Beach, particularly during dry weather summer periods. However, compared to the Santa Ana River, the sewage outfall probably had a negligible impact on surf zone water quality at Huntington Beach and Newport Beach during the storm events sampled in this study. This conclusion is based on the following evidence. First, during our study period, sewage effluent discharged by OCSD was chlorinated and the fecal indicator bacteria concentrations in the final effluent (mean of 6000, 400, and 100 MPN/100 mL for TC, EC, and ENT, $n=17$, C. McGee, personal communication) were significantly below the concentration of fecal indicator bacteria measured in storm water runoff from the Santa Ana River (mean 17000, 5000, and 8000 MPN/100 mL for TC, EC, and ENT, $n=30$, Surbeck et al. (13)). Second, the peak discharge rate from the OCSD outfall (*ca.*, 13 m³/s) is much smaller than the peak discharge rate of storm water runoff from the Santa Ana River (*ca.*, 300 m³/s) (compare blue and black curves, second panel, Figure 2). Third, the sewage effluent is discharged 6 km offshore of the surf zone through a 1 km long diffuser located at the end of OCSD's submarine outfall at a water depth of approximately 60 m (hatched region of the outfall pipe in Figure 1). By contrast, storm water runoff from the Santa Ana River is discharged into the ocean directly at the surf line.

Offshore measurements: satellite ocean color imagery. The spatio-temporal distributions of offshore storm water runoff plumes sampled during this study are revealed by MODIS true color satellite imagery of a 100 km stretch of the coastline centered around our field site (Figure 3). The monitoring grid sampled during the offshore cruises is depicted on the satellite images by yellow dots. The timing of the satellite passes -- relative to rain events, discharge events, wave events, surf zone water quality events, and offshore sampling cruises -- is indicated at the top of Figure 2.

Generally speaking, in this collection of true color imagery the storm water runoff plumes appear to be characterized by a band of turbid water turquoise to brown in appearance that is observed along the entire imaged region, although both cross-shelf and along-shore gradients in the color signature are evident. Following the rain events on 21 - 22 February (total of 39.4 mm, see RE₁ and RE₂ in Figure 2), a MODIS Aqua imagery from 23 February demonstrates the cross-shelf extent of the runoff plume to be variable, ranging from under 1 km in some places to more than 10 km offshore of the Los Angeles River and San Gabriel River (Figure 3A). At our study site, which is centrally located within this broad region, a distinct and apparently heavily particulate-laden runoff plume was observed in the vicinity of the Santa Ana River outlet and nearby station 2201 (see Figure 1 for numerical designation of offshore sampling sites). The Santa Ana River plume extended offshore past station 2203, with an apparent turn down-coast (*i.e.*, southeast), continuing past stations 2104 and 2024. During this time breaking waves were out of the south and the transport direction of fecal indicator bacteria in the surf zone was directed up-coast -- opposite the apparent transport direction of storm water plumes offshore of the surf zone (compare timing of satellite image 1 with WE₂ and fecal indicator bacteria events TC₂, FC₂, and ENT₂, Figure 2). It also appears that a portion of the Los Angeles River and San Gabriel River storm water plumes may have advected south and co-mingled with the Santa Ana River storm water plume. Further south, offshore particulate loadings off the Newport Bay outlet (station 2001) do not appear to be as large as those off the Santa Ana River outlet.





Figure 2. Time series measurements of rainfall, stream discharge at the Santa Ana River and San Gabriel River, and discharge of treated sewage from the OCSD outfall (top panel); water level, salinity, temperature, and chlorophyll measured at the NEOCO sensor (second and third panels); the direction and height of breaking waves at the Newport Beach Pier (fourth panel); and the concentration of fecal indicator bacteria in the surf zone (color contour plots, fifth through seventh panels). Shown at the top of the figure is the timing of the satellite images (blue lettering) and the offshore sampling cruises (black squares).

A MODIS image on 27 February revealed two distinct plumes of considerable size and offshore extent (Figure 3B). This satellite acquisition preceded by one day the sampling cruise on 28 February (described in the next section), followed the large precipitation event on 25 - 26 February (total of 51.3 mm, see RE₃ in Figure 2), and followed the large discharge event from the Santa Ana River (DE₄, in Figure 2). The plume to the northwest in this image appears to be associated with the Los Angeles River and/or San Gabriel River outlets, with an approximate areal extent of 450 km². The plume to the southeast appears to be distinct from the former plume and likely originated from the Santa Ana River outlet, with an approximate areal extent of 100 km² (the presumptive Los Angeles River, San Gabriel River, and Santa Ana River plumes are delineated by red lines in Figure 3B). The 27 February Santa Ana River storm water plume is considerably larger in size than the one observed on 23 February (compare Figures 3A and 3B), consistent with the very large volume of water discharged from the Santa Ana River just prior to this satellite acquisition (approximately 4×10^7 m³, see DE₄ in Figure 2). Further, the Los Angeles River, San Gabriel River, and Santa Ana River runoff plumes on 27 February differed from those on 23 February in that they penetrated farther offshore (30 km compared to 7 km), and thus potentially transported more sediments into the deep waters of the San Pedro Channel.


The jet like appearance of the presumptive Los Angeles River, San Gabriel River, and Santa Ana River storm water runoff plumes in Figure 3B has been observed elsewhere in the Southern California Bight -- e.g., off the Santa Clara River discharge (4, 29) -- and is potentially the result of inertia-driven flow. At the time of this second satellite acquisition breaking waves were out of the west, and along-shore transport in the surf zone, and offshore of the surf zone, appear to be directed down-coast (compare timing of satellite image 2 with WE₃ and fecal indicator events TC₃, FC₃, and ENT₃).

Subsequent MODIS true color imagery on 28 February (Figure 3C) and 29 February (Figure 3D) indicates that both the Los Angeles River/San Gabriel River and Santa Ana River runoff plumes had significantly decreased in size, consistent with reduced flow out of the respective rivers (compare stream discharge curves with timing of satellite images 2 and 3, Figure 2). However, particulate matter appeared to remain high in the general vicinity of the Santa Ana River outlet. Whereas this zone of elevated particulate matter extended south to at least station 2021 on 27 - 28 February, by 29 February it had receded somewhat and was fairly localized around station 2201. Unfortunately no satellite imagery was available the following day (1 March) to complement the third sampling cruise, given persistent regional cloud cover that day.

(A) 23 Feb. at 13:00



(B) 27 Feb. at 12:35



(C) 28 Feb. at 13:20



(D) 29 Feb. at 10:50




Figure 3. MODIS Terra and Aqua true color satellite imagery of storm water runoff plumes along the San Pedro Shelf, California with nominal spatial resolution of 250 m. Yellow dots indicate location of field sampling stations offshore of Huntington and Newport Beach; black arrows denote the Los Angeles River (LAR) outlet, San Gabriel River (SGR) outlet, Santa Ana River/Talbert Marsh (SAR/TM) outlet, and Newport Bay outlet. (A) MODIS-Aqua, 23 February 2004 at 21:00 UTC (13:00), (B) MODIS-Aqua, 27 February 2004 at 20:35 UTC (12:35 local time), (C) MODIS-Aqua, 28 February 2004 at 21:20 UTC (13:20 local time), (D) MODIS-Terra, 29 February 2004 at 18:50 UTC (10:50 local time).

Offshore measurements: *in situ* turbidity and number-averaged particle size. In situ turbidity measurements collected during the three offshore cruises are presented as a series of color contour plots in Figure 4. During the 23 February cruise, a region of high turbidity -- as evidenced by low transmissivity and high TNC -- is evident offshore of, and to the south of, the Santa Ana River outlet (left-hand column of panels, Figure 4). The number-averaged particle size is depressed in this same region, as well as in the region offshore of the Newport Bay outlet. During subsequent cruises, the ocean became progressively less turbid closer to shore (although not necessarily offshore) -- as evidenced by increasing transmissivity and decreasing TNC -- and the number-averaged particle size progressively increased (second and third columns, Figure 4). These results suggest that, offshore of the surf zone, particle size was steadily increasing and particle concentrations were steadily decreasing following the rain and stream discharge events that ended on, or before, the evening of 27 February. The above turbidity patterns are generally

consistent with the plume signatures and gradients observed in the true color satellite imagery (Figure 3), although some differences exist which could result from the offset timing (up to several hours) between the acquisition of the satellite images and the field measurements. As a technical aside, the number-averaged particle size (\bar{d} , see eq 1b) and median particle size (d_{50}) follow similar trends (i.e., they both rise and fall together), although the magnitude of d_{50} was approximately 16 fold larger (Figure S3, Supporting Information). For the results presented here, \bar{d} was chosen because it emphasizes changes in the small end of particle size spectra.




Figure 4. Particle measurements collected during the three sampling cruises. The bottom row of panels indicates the sampling track. TNC is an abbreviation for total particle number concentration. TNC and number-averaged particle size were calculated from measured particle size spectra using eq 1a, b.

Offshore measurements: fecal indicator bacteria. Water quality test results from the three offshore cruises are presented as a set of color contour plots in Figure 5. During the 23 February cruise, the concentration of fecal indicator bacteria exceeded the California single-sample standards for TC, ENT, and EC in several samples collected just offshore, and to the south, of the Santa Ana River and Newport Bay outlets (left-hand column of panels in Figure 5). Nevertheless, the highest concentrations measured offshore of the surf zone are generally lower, in many cases by several orders of magnitude, compared to the highest concentrations measured in the surf zone (compare concentration scales for EC, FC and ENT in Figures 2 and 5). The

difference in offshore and surf zone fecal indicator bacteria concentrations is even more pronounced during the later cruise dates. For example, none of the samples collected during the 28 February and 1 March cruises exceeded State standards for fecal indicator bacteria, yet several of the samples collected from the surf zone during the same time period exceeded single-sample standards for one or more fecal indicator bacteria groups (compare concentrations measured during the second cruise date with TC₃, FC₃, and ENT₃, and concentrations measured during the third cruise date with TC₄, FC₄, and ENT₄, Figures 2 and 5).

Offshore measurements: *F⁺ coliphage and human viruses*. Offshore samples tested positive for *F⁺ coliphage* ($n = 8$, see Table 1), with the exception of a single sample collected on the 28 February cruise from offshore of the Newport Pier (blue, green, and red plus symbols, bottom panels, Figure 5). Human adenoviruses and enteroviruses were detected by real time Q-PCR, nested PCR, and RT-PCR in a sample collected from station 2201 located directly offshore of the Santa Ana River outlet during the 28 February cruise (red plus, middle bottom panel, Figure 5). The concentration of human adenoviruses in this sample is estimated to be 9.5×10^3 genomes per liter of water, which is approximately equivalent to 10 plaque forming units per liter of water, according to a laboratory study comparing Q-PCR results with plaque assay (35). Human enteroviruses were also detected in a sample collected directly offshore of the Santa Ana River outlet (station 2201) on the 23 February cruise (green plus, bottom left panel, Figure 5). While relatively few samples were tested for human viruses ($n = 8$), these results demonstrate that human viruses are present in surface water offshore of the Santa Ana River outlet following storm events, even when the fecal indicator bacteria concentrations are below State standards (e.g., station 2201 during the 28 February cruise, Figure 5). These results are consistent with previous observations that human pathogenic viruses and fecal indicator viruses persist longer than fecal indicator bacteria in ocean water (36). It should also be noted that direct PCR measurement of pathogenic viruses in highly turbid water is challenging due to PCR inhibition (35).




Figure 5. Fecal indicator bacteria concentrations measured during the three sampling cruises. The bottom row of panels indicates the sampling track (blue arrows), and the detection of F⁺ coliphage and human viruses. SAR/TM is an abbreviation for the outlet of the Santa Ana River and Talbert Marsh.

Offshore measurements: Relationship between fecal indicator bacteria, turbidity, and number-averaged particle size. Turbidity has been suggested as a possible proxy for water quality (37,38). However, based on our offshore data, turbidity per se appears to be an inconsistent proxy for the concentration of fecal indicator bacteria. For example, during the 23 February cruise there is good coherence between turbidity and TC, EC and ENT concentrations off the Santa Ana River outlet and Newport Pier (compare transmissivity and TNC with fecal indicator bacteria results, left-hand column of panels, Figures 4 and 5). However, turbidity is low off of the Newport Bay outlet where the bacteria concentrations are particularly high. In addition, there are no consistently robust relationships between shipboard measurements of fecal indicator bacteria and shipboard measurements of TOC, temperature, or salinity (see Figure S4, Supporting Information). The number-averaged particle size, on the other hand, comes close to matching the along-shore spatial pattern of fecal indicator bacteria measured during the 23 February cruise. Specifically, elevated fecal indicator bacteria concentration appears to correlate with depressed number-averaged particle size (compare fecal indicator bacteria and number-averaged particle size results for the 23 February cruise, left-hand column of panels, Figures 4

and 5). When all of the fecal indicator bacteria data collected during the three cruises are aggregated and plotted against number-averaged particle size, an inverse relationship between these two parameters emerges; specifically, samples with elevated fecal indicator bacteria concentrations also exhibit small number-averaged particle size (Figure 6A). Moreover, the concentration of fecal indicator bacteria in water samples collected during the first two cruises is the same, within error, before and after filtration through a 53 μm sieve (Figure 6B), implying that fecal indicator bacteria are either adsorbed to particles smaller than 53 μm , or are not particle-associated. TOC also appears to pass through the 53 μm sieve (Figure 6B) as do human viruses and fecal indicator viruses (data not shown). The co-occurrence of small particles and indicators of fecal pollution (fecal indicator bacteria, fecal indicator viruses, and human pathogenic viruses) does not necessarily imply that the latter are adsorbed to the former. The inverse relationship evident in Figure 6A, for example, may reflect a temporal evolution of storm water plumes as they age, from a predominance of small particles and high concentrations of fecal indicators initially, to larger particles and lower concentrations of fecal indicators later.




Figure 6. (A) Cross plots of log-transformed fecal indicator bacteria concentrations measured in samples collected during the three offshore cruises, against the corresponding number-averaged particle size. (B) Cross plots of log-transformed fecal indicator bacteria concentrations and TOC concentrations measured in samples collected during the three offshore cruises, before and after

filtration through a 53 μm sieve. The one-to-one line corresponds to the case where the concentrations are the same before and after filtration.

Offshore measurements: particle size spectra. Particle size spectra acquired during the three cruises are presented in Figure 7. Each plot displays the normalized particle volume (vertical axis) detected in 32 logarithmically spaced particle diameter bins ranging in size from 2.5 to 500 μm (horizontal axis). The particle size spectrum measured at a particular offshore location and time appear to be related to the specific storm water plume the particles are associated with and, possibly the elapsed time storm water has spent in the ocean. Storm water flowing out of the Santa Ana River during the 23 February cruise, for example, is characterized by two modes at the small end of the size spectrum, one in the $< 5 \mu\text{m}$ bin and another in the 10 to 50 μm bins (set of red curves, Figure 7). These modes are present in storm water runoff sampled at several locations in the Santa Ana River watershed (13), in samples collected at the ocean outlet of the Santa Ana River (panel labeled “SAR Outlet” at top of Figure 7), and in samples collected just offshore (red curve at station 2201, Figure 7) and down-coast (red curve at station 2101, Figure 7) of the Santa Ana River outlet. Particles discharged from the Santa Ana River appear to dilute and merge into a background turbidity characterized by a single broad mode in the 50 to 300 μm size range (evident in the red curves at most stations, Figure 7).

Referring to Figure 3A and the earlier discussion of this satellite image, the 50 to 300 μm mode observed on 23 February may be characteristic of a large runoff plume originating from one or more up-coast sources of storm water runoff, most likely the Los Angeles River and/or the San Gabriel River. It should be noted that a number of factors can lead to artifacts in the particle size spectra estimated from the light scattering instrument deployed in this study (39). However, in our case this caveat is mitigated somewhat by the observation that particle volume fractions calculated from the particle size spectra are strongly correlated (Spearman’s rank correlation $Sp = 0.90$, $p = 0.02$) with independent measurements of total suspended solids (data not shown).

During the second and third cruises, the particle size spectra progressively coarsen with the result that, by 1 March, virtually all of the particle volume is associated with the largest size bin ($> 500 \mu\text{m}$, green curves in Figure 7). The observed temporal evolution in particle size spectra -- from high turbidity and multiple modes at the lower end of the particle size spectrum to low turbidity and a single mode at the large end of the particle size spectrum -- may reflect decreasing particle supply (i.e., reduced storm water discharge from major river outlets) coupled with within-plume coagulation of particles into larger size classes and, ultimately, removal of the largest particles by gravitational sedimentation. Coagulation time scales estimated from these particle size spectra measurements are short (minutes to hours or longer) compared to time scales associated with the generation and offshore transport of storm water plumes (hours to days), and hence coagulation cannot be ruled out as an important mechanism at our field site (see Supporting Information for details on the time scale calculations). Whether coagulation, in fact, plays a role in the fate and transport of particles and particle-associated contaminants in storm water plumes will likely depend on the coagulation efficiency (i.e., the fraction of particle-particle collisions that result in sticking events) and shear rates present at a given location and time (40, 41). Alternatively, the observed temporal coarsening of particles in the offshore may reflect changes in the particle size spectra of the storm water runoff before it enters the ocean, from a predominance of smaller particles during the peak of the hydrograph, to a predominance of coarser particles during the falling limb of the hydrograph. Further studies are needed to determine whether observed coarsening of the offshore particle size spectra is caused by within-

plume coagulation, and/or temporal evolution of the particle size spectra in storm water runoff before it enters the ocean.




Figure 7. Particle size spectra measured during the three offshore cruises; numbers at the top of each panel denote the station number where the particle size spectra were measured (see Figure 1). The vertical axis in each plot represents the particle volume resident in logarithmically spaced particle diameter bins; the horizontal axis represents the diameter of the particles (in μm). These plots are arranged so that the stations progress from onshore to offshore (top to bottom) and up-coast to down-coast (left to right). The single plot labeled “SAR Outlet” corresponds to a particle size spectrum measured in storm water runoff flowing out of the Santa Ana River outlet, just upstream of where it flows over the beach and into the ocean.

Data Synthesis

Results presented in this paper are represented schematically in Figure 8, including potential offshore transport mechanisms (panel A) and the resulting distribution of particles, bacteria, and viruses (panel B). As storm water is discharged from the river outlet and flows over the beach, a fraction is entrained in the surf zone and the rest is ejected offshore in a momentum jet. Measurements of fecal indicator bacteria in the surf zone suggest that, once entrained, contaminants are transported parallel to shore by wave-driven currents, in a direction (i.e., up or down-coast) controlled by the approaching wave field. When waves strike the beach so that a component of wave momentum is directed up-coast (the scenario pictured in Figure 8), fecal indicator bacteria in the surf zone are carried up-coast of the river outlet. Conversely, when waves strike the beach so that a component of wave momentum is directed down-coast, fecal indicator bacteria in the surf zone are carried down-coast of the river outlet. The build-up of

water in the surf zone from breaking waves drives a cross-shore circulation cell, which can transport material between the surf zone and offshore of the surf zone. At our field site, this cross-shore circulation appears to limit the length of beach severely polluted with fecal indicator bacteria to < 5 km around the river outlet, by diluting contaminated surf zone water with cleaner water from offshore. While the transport processes described here are based on measurements of fecal indicator bacteria in the surf zone, it is likely that other contaminants in storm water runoff -- in particular, human viruses, and toxic contaminants associated with suspended particles (13, 42) -- will behave similarly.


Further offshore, storm water runoff plumes are common and readily detected through a variety of geophysical parameters (e.g., salinity, transmissivity, surface color). A clear linkage between these parameters and fecal indicator bacteria could not be established here. However, fecal indicator bacteria did appear to be associated with the smallest particle sizes, based on both fractionation studies (Figure 6B) and the inverse relationship observed between fecal indicator bacteria concentrations and number-averaged particle size (Figure 6A). Particle size spectra in the offshore plumes coarsen with time post-release, and fecal indicator bacteria concentrations steadily drop (see the schematic representation of particle size in the various offshore plumes, Figure 8B). These results have several implications. First, they suggest that high concentrations of fecal indicator bacteria in the surf zone at our field site are probably not brought into the study area by coastal currents from distal sources (e.g., the Los Angeles river and/or San Gabriel river). Second, cross-shore transport of water between the surf zone and offshore of the surf zone -- for example, by rip cell currents -- is likely to improve surf zone water quality by diluting dirty river effluent entrained in the surf zone with relatively clean ocean water from offshore.

While the concentrations of fecal indicator bacteria in the offshore plumes are generally below surf zone water quality standards, particularly during the latter two cruises, fecal indicator viruses (F^+ coliphage) were detected in nearly all offshore samples tested, and human adenoviruses and enteroviruses were detected in several offshore samples, including two collected offshore of the Santa Ana River outlet (station 2201 on 23 and 28 February, see Figure 5). It is likely that the virus results presented here represent a conservative estimate of viral prevalence, because a limited numbers of samples were tested ($n = 8$). In addition, the presence of PCR inhibitors in storm water reduces the efficiency of PCR detection of human pathogenic viruses, as mentioned earlier. At present, there are no water quality standards for fecal indicator viruses and human pathogenic viruses, largely because epidemiological data are not available to link adverse human health outcomes (e.g., gastrointestinal disease) to recreational ocean exposure to these organisms. However, the offshore detection of human pathogenic viruses begs several questions: First, do these viruses constitute a human health risk, either by contaminating the surf zone direction (see arrow with question mark, indicting the possible transfer of contaminants from offshore into the surf zone, Figure 8(B)), or by sequestering in offshore sediments? Second, given the fact that the Santa Ana River has separate storm and sanitary sewer systems, what is the source of human fecal pathogens in the wet weather water runoff? Many studies have shown that human fecal pathogens are associated with storm runoff from urban areas located throughout the U.S. (25,43-45) -- so the association between storm water runoff and human fecal pathogens observed here is certainly not unique. Possible sources of human pathogens in storm water runoff from urban areas include leaking sewer pipes, illicit sewage connections to the storm water sewer system, homeless populations, etc.

Taken together, the results presented in this paper demonstrate that storm water runoff from the Santa Ana River is a significant source of near shore pollution, including turbidity, fecal indicator bacteria, fecal indicator viruses, and human pathogenic viruses. However, relationships

between variables (e.g., between turbidity and fecal indicator bacteria, and between fecal indicator bacteria and human viruses) vary from site to site (at the same time), and from time to time (at the same site) suggesting that the sources, fate, and transport processes are contaminant specific. The apparent exception is the inverse relationship observed between fecal indicator bacteria and number-averaged particle size, although further studies are needed to determine if this result is generalizable to other storm seasons and coastal sites and, if so, to determine the underlying mechanism at work. The relationship between water quality parameters (e.g., fecal indicator bacteria), turbidity and other field proxies -- such as number-averaged particle size, salinity, colored dissolved organic matter -- are the focus of ongoing and future regional studies, including as part of a coastal water quality observing program within the Bight '03 Project (http://www.sccwrp.org/regional/03bight/bight03_fact_sheet.html), as well as other investigations being carried out as part of the Southern California Coastal Ocean Observing System (SCCOOS).

A)



B)




Figure 8. (A) Transport mechanisms that can affect the offshore distribution of contaminants discharged from river outlets. (B) Schematic representation of the spatial distribution of particles (black circles of varying size), fecal indicator bacteria (red symbols), and F^+ coliphage and human pathogenic viruses (green symbols). Abbreviations are SAR (Santa Ana River), SGR (San Gabriel River) and LAR (Los Angeles River).

Acknowledgements

This study was funded by a joint grant from the National Water Resource Institute (03-WQ-001) and the US Geological Survey National Institutes for Water Research (UCOP-33808), together with matching funds from the Counties of Orange, Riverside, and San Bernardino in southern California. MODIS data were acquired as part of the NASA's Earth Science Enterprise, and processed by the MODIS Adaptive Processing System (MODAPS), the Goddard Distributed Active Archive Center (DAAC), and are archived and distributed by the Goddard DAAC. The JPL effort was supported by the National Aeronautics and Space Administration through a contract with the Jet Propulsion Laboratory, California Institute of Technology. NEOCO measurements were supported by the University of California Marine Council's Coastal Environmental Quality Initiative. Partial support for human virus and fecal indicator virus study was provided by Water Environmental Research Foundation award 01-HHE-2a. We also acknowledge the contribution of Weiping Chu at UCI for technical assistance with human viruses analysis. The authors acknowledge the input and feedback from numerous colleagues, most notably Chris Crompton, George L. Roberson, Charles D. McGee, Rick Wilson, Brett F. Sanders, Patricia Holden, Ronald Linsky, and Steve Weisberg. Sample collection and processing was carried out with the help of Youngsul Jeong and Ryan Reeves. The authors also thank the Assistant Manager of the City of Newport Beach, David Kiff, the Chief of the Newport Beach Fire Department, Timothy Riley, John Moore, and Brian O'Rourke for arranging the 23 February cruise, and officials at the Orange County Sanitation District for assisting in the collection and analysis of offshore and surf zone water samples. Some of the data and ship time for this study were donated by the Bight'03 program. The authors also acknowledge the excellent feedback provided on the manuscript by three anonymous reviewers.

Supporting Information Available.

Sampling and analysis protocols, calculation of the orthokinetic coagulation time scales, and additional figures.

References

- (1) Culliton, T. J. *Population; distribution, density and growth; A state of the coast report*; NOAA's state of the coast report; National Oceanic and Atmospheric Administration: Silver Spring, MD, 1998.
- (2) Reeves, R. L.; Grant, S. B.; Mrse, R. D.; Copil Oancea, C. M.; Sanders, B. F.; Boehm, A. B. Scaling and management of fecal indicator bacteria in runoff from a coastal urban watershed in southern California. *Environ. Sci. Technol.* **2004**, *38*, 2637-2648.
- (3) Bay, S.; Jones, B. H.; Schiff, K.; Washburn L. Water quality impacts of stormwater discharges to Santa Monica Bay. *Mar. Environ. Res.* **2003**, *56*, 205-223.
- (4) Warrick, J. A.; Mertes, L. A. K.; Washburn, L.; Siegel, D. A. Dispersal forcing of southern California river plumes, based on field and remote sensing observations. *Geo-Mar. Lett.* **2004**, *24*, 46-52.

- (5) Koh, R. C. Y.; Brooks, N. H. Fluid mechanics of wastewater disposal in the ocean. *Annual Rev. Fluid Mech.* **1975**, 7, 187-211.
- (6) Lu, R.; Turco, R. P.; Stolzenbach, K.; Fiedlander, S. K.; Xiong, C. Dry deposition of airborne trace metals on the Los Angeles Basin and adjacent coastal waters. *J. Geophys. Res.-Atmos.* **2003**, 108, AAC 11, 1-24.
- (7) Boehm, A. B.; Shellenbarger, G. G.; Paytan, A. Groundwater discharge: potential association with fecal indicator bacteria in the surf zone. *Environ. Sci. Technol.* **2004**, 38, 3558-3566.
- (8) Schiff, K.C. *Development of a model publicly owned treatment work (POTW) monitoring program*; Southern California Coastal Water Research Project: Westminster, CA, 1999.
- (9) Warrick, J. A.; Rubin, D. M.; Orzech K. M. The effects of urbanization and flood control on suspended sediment discharge of a southern California river, evidence of a dilution effect. *Water Resources Res.* **2004**, submitted.
- (10) Schiff, K. C.; Allen, M. J.; Zeng, E. Y.; Bay, S. M. Southern California. *Mar. Pollut. Bull.* **2000**, 41, 76-93.
- (11) King, J. A.; Leubs, R. A.; Hardy, W. T.; Smith, A. B.; Withers, J. B.; Reynolds, A.; Henriques, M.; Johnson, T.; Thibeault, G. J. *Water quality control plan; Santa Ana River Basin* (8); California Regional Water Quality Control Board: Santa Ana Region, 1995.
- (12) DiGiacomo, P. M.; Washburn L.; Holt, B.; Jones, B. H. Coastal pollution hazards in southern California observed by SAR imagery: stormwater plumes, wastewater plumes, and natural hydrocarbon seeps. *Mar. Pollut. Bull.* **2004**, 49, 1013-1024.
- (13) Surbeck, C. Q.; Grant, S. B.; Ahn, J. H.; Jiang, S. Transport of suspended particles and fecal pollution in storm water runoff from an urban watershed in southern California. *Environ. Sci. Technol.* **2004**, submitted.
- (14) Grant, S. B.; Sanders, B. F.; Boehm, A. B.; Redman, J. A.; Kim, J. H.; Mrse, R. D.; Chu, A. K.; Gouldin, M.; McGee, C. D.; Gardiner, N. A.; Jones, B. H.; Svejksky, J.; Leipzig, G. V.; Brown, A. Generation of Enterococci Bacteria in a coastal saltwater marsh and its impact on surf zone water quality. *Environ. Sci. Technol.* **2001**, 35, 2407-2416.
- (15) Boehm, A. B.; Grant, S. B.; Kim, J. H.; Mowbray, S. L.; McGee, C. D.; Clark, C. D.; Foley, D. M.; Wellman, D. E. Decadal and shorter period variability of surf zone water quality at Huntington Beach, California. *Environ. Sci. Technol.* **2002**, 36, 3885-3892.
- (16) Boehm, A. B.; Sanders, B. F.; Winant, C. D. Cross-shelf transport at Huntington Beach: Implications for the fate of sewage discharged through an offshore ocean outfall. *Environ. Sci. Technol.* **2002**, 36, 1899-1906.
- (17) Grant, S. B.; Sanders, B. F.; Boehm, A. B.; Arega F.; Ensari, S.; Mrse, R. D.; Kang H. Y.; Reeves R. L.; Kim, J. H.; Redman, J. A. *Coastal runoff impact study phase II: Sources and*

dynamics of fecal indicators in the lower Santa Ana River Watershed; A draft report prepared for the National Water Research Institute: County of Orange, and the Santa Ana Regional Water Quality Control Board, 2002.

- (18) Turbow, D.; Lin, T. H.; Jiang, S. Impacts of beach closures on perceptions of swimming-related health risk in Orange County, California. *Mar. Pollut. Bulletin* **2004**, 48, 132-136.
- (19) Jones, B. H.; Noble, M. A.; Dickey, T. D. Hydrographic and particle distributions over the Palos Verdes Continental Shelf: spatial, seasonal and daily variability. *Cont. Shelf Res.* **2002**, 22, 945-965.
- (20) Washburn, L.; McClure, K. A.; Jones, B. H.; Bay, S. M. Spatial scales and evolution of stormwater plumes in Santa Monica Bay. *Mar. Environ. Res.* **2003**, 56, 103-125.
- (21) DeLeon, R.; Shieh, Y. S. C.; Baric, R. S.; Sobey, M. D. Detection of enteroviruses and hepatitis A virus in environmental samples by gene probes and polymerase chain reaction. *Water Quality Conference*; American Water Works Association: Denver, CO, 1990, 833-853.
- (22) Tsai, Y. L.; Sobsey, M. D.; Sangermano, L. R.; Palmer, C. J. Simple method of concentrating enteroviruses and hepatitis A virus from sewage and ocean water for rapid detection by reverse transcriptase-polymerase chain reaction. *Appl. Environ. Microbiol.* **1993**, 59, 3488-3491.
- (23) Jiang, S. C.; Chu, W. PCR detection of pathogenic viruses in southern California urban river. *J. Appl. Microbiol.* **2004**, 97, 17-28.
- (24) Pina, S.; Puig, M.; Lucena, F.; Jofre, J.; Girones, R. Viral pollution in the environment and in shellfish: Human adenovirus detection by PCR as an index of human viruses. *Appl. Environ. Microbiol.* **1998**, 64, 3376-3382.
- (25) He, J.; Jiang, S. Quantification of enterococci and human adenoviruses in environmental samples by real-time PCR. *Appl. Environ. Microbiol.* **2004**, *in press*.
- (26) Mikkelsen, O. A. Variation in the projected surface of suspended particles: Implications for remote sensing assessment of TSM. *Rem. Sens. of Environ.* **2002**, 79, 23-29.
- (27) Serra, T.; Colmer, J.; Cristina, X. P.; Vila, X.; Arellano, J. B.; Casamitjana, X. J. Evaluation of laser in-situ instrument for measuring concentration of phytoplankton, purple sulfur bacteria, and suspended inorganic sediments in lakes. *Environ. Eng.* **2001**, 11, 1023-1030.
- (28) Kim, J. H.; Grant, S. B.; McGee, C. D.; Sanders, B. F.; Largier, J. L. Locating sources of surf zone pollution: A mass budget analysis of fecal indicator bacteria at Huntington Beach, California. *Environ. Sci. Technol.* **2004**, 38, 2626-2636.
- (29) Inman, D. L.; Brush, B. M. Coastal challenge. *Science*. **1973**, 181, 20-32.

- (30) AES Huntington Beach generating station surf zone water quality study final draft; A consultant report prepared for California Energy Commission; KOMEX H2O Science Incorporated: Westminster, CA, 1998.
- (31) Boehm, A. B. Model of microbial transport and inactivation in the surf zone and application to field measurements of total coliform in northern Orange County, California. *Environ. Sci. Technol.* **2003**, 37, 5511-5517.
- (32) Boehm, A. B.; Sanders, B. F.; Winant, C. D. Cross-shelf transport at Huntington Beach. Implications for the fate of sewage discharged through an offshore ocean outfall. *Environ. Sci. Technol.* **2002**, 36, 1899-1906.
- (33) Boehm, A. B.; Lluch-Cota D. B.; David, K. A.; Winant, C. D.; Monismith S. G. Covariation of coastal water temperature and microbial pollution at interannual to tidal periods. *Geophys. Res. Lett.* **2004**, 31, L06309.
- (34) Mertes, L. A. K.; Warrick, J. A. Measuring flood output from 110 coastal watersheds in California with field measurements and SeaWiFS. *Geology* **2001**, 29, 659-662.
- (35) Jiang, S. C.; Deszfulian, H.; Chu, W. Real-time quantitative PCR for enteric adenovirus serotype 40 in environmental waters. *Can. J. Microbiol.* **2004**, submitted.
- (36) Shuval, H. I. *Developments in Water Quality Research*; Ann Arbor-Humphrey Science: Ann Arbor, 1970.
- (37) Boucier, D. R.; Sharma, R. P. Heavy metals and their relationship to solids in urban runoff. *Int. J. Envir. Anal. Chem.* **1980**, 7, 273-283.
- (38) Gippel, C. J. Potential of turbidity monitoring for measuring the transport of suspended-solids in streams. *Hydrological Processes.* **1995**, 9, 83-97.
- (39) Mikkelsen O. A. In-situ particle size spectra and density of particle aggregates in a dredging plume. *Mar. Geo.* **2000**, 170, 443-459.
- (40) Grant, S. B.; Poor, C.; Relle, S. Scaling theory and solutions for the steady-state coagulation and settling of fractal aggregates in aquatic systems. *Colloid and Surfaces.* **1996**, 107, 155-174.
- (41) Grant, S. B.; Kim, J. H.; Poor, C. Kinetic theories for the coagulation and sedimentation of particles. *J. Colloid Interfaces Sci.* **2001**, 238, 238-250.
- (42) Glenn, D. W.; Sansalone, J. J. Accretion and partitioning of heavy metals associated with snow exposed to urban traffic and winter storm maintenance activities. II. *J. Environ. Eng. ASCE.* **2002**, 2, 167-185.
- (43) Lipp, E. K.; Kurz, R.; Vincent, R.; Rodriguez-Palacios, C.; Farrah, S. R.; Rose, J. R.; The effects of seasonal variability and weather on microbial fecal pollution and enteric pathogens in a subtropical estuary. *Estuaries.* **2001**, 24, 266-276.

(44) Noble, R. T.; Fuhrman, J. A.; Enteroviruses detected by reverse transcriptase polymerase chain reaction from the coastal waters of Santa Monica Bay, California: low correlation to bacterial indicator levels. *Hydrobiologia*. **2001**, *460*, 175-184.

(45) Jiang, S. C.; Chu., W.; PCR detection of pathogenic viruses in southern California urban rivers. *J. App. Microbiol.* **2004**, *97*, 17-28.

Synopsis

The coastal water quality impact of storm water runoff from an urban watershed in southern California depends on prevailing ocean currents, within-plume processing of particles and pathogens, and the timing, magnitude and nature of runoff discharged from river outlets over the course of a storm.