

SCCWRP #0637

## Plastic ingestion by planktivorous fishes in the North Pacific Gyre

Christiana M. Boerger<sup>1</sup>, Gwendolyn L. Lattin<sup>1</sup>, Shelly L. Moore<sup>2</sup> and Charles J. Moore<sup>1</sup>

<sup>1</sup>*Algalita Marine Research Foundation, Long Beach, CA*

<sup>2</sup>*Southern California Coastal Water Research Project, Costa Mesa, CA*

### ABSTRACT

A significant amount of marine debris has accumulated in the North Pacific Central Gyre (NPCG). The effects on larger marine organisms have been documented through cases of entanglement and ingestion; however, little is known about the effects on lower trophic level marine organisms. This study is the first to document ingestion and quantify the amount of plastic found in the gut of common planktivorous fish in the NPCG. From February 11 to 14, 2008, 11 neuston samples were collected by manta trawl in the NPCG. Plastic from each trawl and fish stomach was counted and weighed and categorized by type, size class and color. Approximately 35% of the fish studied had ingested plastic, averaging 2.1 pieces per fish. Additional studies are needed to determine the residence time of ingested plastics and their effects on fish health and the food chain implications.

**Due to distribution restrictions, the full-text version of this article is available by request only.**

Please contact [pubrequest@sccwrp.org](mailto:pubrequest@sccwrp.org) to request a copy.